

CD11b Antibody

Rabbit mAb

Catalog # AP90086

Product Information

Application	WB, IHC, IF, ICC, IHF
Primary Accession	P11215
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	CR3A; MO1A; CD11B; MAC-1; MAC1A; SLEB6;ITGAM
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	127179

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human CD11b
Description	ITGAM, also named as CD11B and CR3A, belongs to the integrin alpha chain family. It is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles. ITGAM is identical with CR-3, the receptor for the iC3b fragment of the third complement component. It probably recognizes the R-G-D peptide in C3b.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	ITGAM (HGNC:6149)
Synonyms	CD11B, CR3A
Function	Integrin ITGAM/ITGB2 is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles and pathogens (PubMed: 20008295 , PubMed: 9558116). It is identical with CR-3, the receptor for the iC3b fragment of the third complement component. It probably recognizes the R-G-D peptide in C3b. Integrin ITGAM/ITGB2 is also a receptor for fibrinogen and factor X. It recognizes P1 and P2 peptides of fibrinogen gamma chain. Regulates neutrophil migration (PubMed: 28807980). In association with beta subunit ITGB2/CD18, required for CD177-PRTN3-mediated activation of TNF primed neutrophils (PubMed: 21193407). Integrin ITGAM/ITGB2 is also a receptor for ICAM1 ligand ensuring adhesion between stimulated neutrophils

and stimulated endothelial cells (PubMed:[1980124](#)). May regulate phagocytosis-induced apoptosis in extravasated neutrophils (By similarity). May play a role in mast cell development (By similarity). Required with TYROBP/DAP12 in microglia to control production of microglial superoxide ions which promote the neuronal apoptosis that occurs during brain development (By similarity).


Cellular Location

Cell membrane; Single-pass type I membrane protein. Membrane raft; Single-pass type I membrane protein

Tissue Location

Predominantly expressed in monocytes and granulocytes (PubMed:[1346576](#)). Expressed in neutrophils (at protein level) (PubMed:[21193407](#)).

Images



Western blot analysis of CD11b expression in RAW264.7 cell lysate.

Image not found : 202311/AP90086-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human spleen, using CD11b Antibody.

Image not found : 202311/AP90086-wb6.jpg

Zinc inhibited LPS-induced inflammatory responses by upregulating A20 expression in microglia BV2 cells. -Journal of Affective Disorders

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.