

5T4 Antibody

Rabbit mAb Catalog # AP90363

Product Information

Application WB, IP
Primary Accession Q13641

Reactivity Rat, Human, Mouse

Clonality Monoclonal

Other Names 5T4; 5T4AG; M6P1; TPBG; 5T4 oncofetal antigen; Trophoblast glycoprotein;

IsotypeRabbit IgGHostRabbitCalculated MW46032

Additional Information

Dilution WB 1:500~1:2000 IP 1:50 **Purification** Affinity-chromatography

Immunogen A synthesized peptide derived from human 5T4

Description 5T4 is a 72 kDa transmembrane glycoprotein that exhibits restricted

expression in human and mouse adult tissues, is upregulated on many carcinomas and tumour expression correlates with poorer clinical outcome in ovarian, gastric and colorectal cancers. mES cells lack cell surface expression of the 5T4 antigen, and both protein and mRNA are rapidly upregulated following induction of differentiation. This proteins expression correlates with

downregulation of OCT-4 and Tra-1-60.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name TPBG

Synonyms 5T4

Function May function as an inhibitor of Wnt/beta-catenin signaling by indirectly

interacting with LRP6 and blocking Wnt3a-dependent LRP6 internalization.


Cellular Location Cell membrane; Single-pass type I membrane protein

Tissue Location Expressed by all types of trophoblasts as early as 9 weeks of development.

Specific for trophoblastic cells except for amniotic epithelium. In adult tissues, the expression is limited to a few epithelial cell types but is found on a variety

of carcinoma

Images



Western blot analysis of 5T4 expression in MCF-7 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.