

Histone H2A (hydroxyl Y39) Antibody

Rabbit mAb

Catalog # AP90477

Product Information

Application	WB, IHC
Primary Accession	P04908
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	H2A.1; H2A/c; H2A1; H2AFC; H2AFD; H2AFI; H2AFN; H2AFP; HIST1H2AG; HIST1H2AI; HIST1H2AK; HIST1H2AL; HIST1H2AM; histone cluster 1, H2ai; Histone H2A type 1; Histone H2A/p;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	14135


Additional Information

Dilution	WB 1:500~1:2000 IHC 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Histone H2A (hydroxyl Y39)
Description	H2A.1 Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	H2AC4 (HGNC:4734)
Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Cellular Location	Nucleus. Chromosome.

Images



Western blot analysis of Calreticulin expression in (1) NIH/3T3 cell lysate; (2) A549 cell lysate.

Image not found : 202311/AP90477-IHC.jpg

Immunohistochemical analysis of paraffin-embedded mouse colon, using Histone H2A (hydroxyl Y39) Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.