

CD3 epsilon Antibody

Rabbit mAb

Catalog # AP90584

Product Information

| | |
|--------------------------|-------------------------|
| Application | WB, IHC, FC, IP |
| Primary Accession | P07766 |
| Reactivity | Human |
| Clonality | Monoclonal |
| Other Names | T3E; TCRE; IMD18; CD3E; |
| Isotype | Rabbit IgG |
| Host | Rabbit |
| Calculated MW | 23147 |

Additional Information

| | |
|-------------------------------------|--|
| Dilution | WB 1:500~1:2000 IHC 1:50~1:200 IP 1:30 FC 1:100 |
| Purification | Affinity-chromatography |
| Immunogen | A synthesized peptide derived from human CD3 epsilon |
| Description | The protein encoded by this gene is the CD3-epsilon polypeptide, which together with CD3-gamma, -delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T-cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

Protein Information


| | |
|-----------------|--|
| Name | CD3E |
| Synonyms | T3E |
| Function | Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response (PubMed: 15294938 , PubMed: 15546002 , PubMed: 2470098 , PubMed: 40592325 , PubMed: 8490660). When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD247/CD3Z (PubMed: 2470098 , PubMed: 40592325). All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain (PubMed: 2470098 , PubMed: 40592325). Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting |

in the activation of downstream signaling pathways (PubMed:[2470098](#), PubMed:[40592325](#)). CD3E ITAM phosphorylation creates docking sites for the protein kinase ZAP70 leading to ZAP70 phosphorylation and its conversion into a catalytically active enzyme (By similarity). In addition of this role of signal transduction in T-cell activation, CD3E plays an essential role in correct T-cell development (By similarity). Also participates in internalization and cell surface down-regulation of TCR-CD3 complexes via endocytosis sequences present in CD3E cytosolic region (PubMed:[10384095](#), PubMed:[26507128](#)). In addition to its role as a TCR coreceptor, it serves as a receptor for ITPR1L1 (PubMed:[38614099](#)). Ligand recognition inhibits T-cell activation by promoting interaction with NCK1, which prevents CD3E-ZAP70 interaction and blocks the ERK- NFkB signaling cascade and calcium influx (PubMed:[12110186](#), PubMed:[38614099](#)).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Images



Western blot analysis of CD3 epsilon expression in Jurkat cell lysate.

Image not found : 202311/AP90584-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human colon cancer, using CD3 epsilon Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.