

Dnmt1 Antibody

Rabbit mAb

Catalog # AP91080

Product Information

Application	WB, IHC, IF, ICC, IHF
Primary Accession	P26358
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	ADCADN; CXXC finger protein 9; CXXC9; DNA methyltransferase 1; DNA MTase; Dnmt1o; HSN1E; M.HsaI; MCMT; Met1; MommeD2;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	183165

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Dnmt1
Description	Methylation of DNA at cytosine residues in mammalian cells is a heritable, epigenetic modification that is critical for proper regulation of gene expression, genomic imprinting and development. It is responsible for maintaining methylation patterns established in development. DNA methylation is coordinated with methylation of histones. Mediates transcriptional repression by direct binding to HDAC2.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	DNMT1 {ECO:0000303 Ref.3, ECO:0000312 HGNC:HGNC:2976}
Function	DNA methyltransferase that methylates CpG residues (PubMed: 17200670 , PubMed: 18754681 , PubMed: 21745816 , PubMed: 26070743). Preferentially methylates hemimethylated DNA (PubMed: 21745816 , PubMed: 26070743). Associates with DNA replication sites in S phase maintaining the methylation pattern in the newly synthesized strand, that is essential for epigenetic inheritance (PubMed: 17200670 , PubMed: 21745816). Associates with chromatin during G2 and M phases to maintain DNA methylation independently of replication (PubMed: 21745816). It is responsible for maintaining methylation patterns established in development (PubMed: 21745816). DNA methylation is coordinated with methylation of histones (PubMed: 16357870). Mediates transcriptional repression by direct binding to HDAC2 (PubMed: 10888872). In association with DNMT3B and via

the recruitment of CTCFL/BORIS, involved in activation of BAG1 gene expression by modulating dimethylation of promoter histone H3 at H3K4 and H3K9 (PubMed:[18413740](#)). Probably forms a corepressor complex required for activated KRAS-mediated promoter hypermethylation and transcriptional silencing of tumor suppressor genes (TSGs) or other tumor-related genes in colorectal cancer (CRC) cells (PubMed:[24623306](#)). Also required to maintain a transcriptionally repressive state of genes in undifferentiated embryonic stem cells (ESCs) (PubMed:[24623306](#)). Associates at promoter regions of tumor suppressor genes (TSGs) leading to their gene silencing (PubMed:[24623306](#)).


Cellular Location

Nucleus. Chromosome Note=Associates with replication foci during S-phase: recruited to hemimethylated DNA sites via its RFTS domain, which specifically recognizes and binds histone H3 ubiquitinated at 'Lys-14', 'Lys-18' and 'Lys-23' (H3K14ub, H3K18ub and H3K23ub, respectively) (PubMed:29053958). Localized to the perinucleolar region (PubMed:24492612).

Tissue Location

Ubiquitous; highly expressed in fetal tissues, heart, kidney, placenta, peripheral blood mononuclear cells, and expressed at lower levels in spleen, lung, brain, small intestine, colon, liver, and skeletal muscle. Isoform 2 is less expressed than isoform 1.

Images



Western blot analysis of Dnmt1 expression in (1) HEK293 cell lysate; (2) NIH/3T3 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.