

IL10 Antibody

Rabbit mAb

Catalog # AP91136

Product Information

Application	WB, IF, FC, ICC
Primary Accession	P22301
Reactivity	Human, Mouse
Clonality	Monoclonal
Other Names	IL10; CSIF; GVHDS; IL10A; TGIF; Interleukin-10;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	20517

Additional Information

Dilution	WB 1:500~1:2000 ICC/IF 1:50~1:200 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human IL10
Description	Interleukin-10 (IL-10) is an anti-inflammatory cytokine that is produced by T cells, NK cells, and macrophages. IL-10 initiates signal transduction by binding to a cell surface receptor complex consisting of IL-10 RI and IL-10 RII, leading to the activation of Jak1 and Tyk2 and phosphorylation of Stat3.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	IL10
Function	Major immune regulatory cytokine that acts on many cells of the immune system where it has profound anti-inflammatory functions, limiting excessive tissue disruption caused by inflammation. Mechanistically, IL10 binds to its heterotetrameric receptor comprising IL10RA and IL10RB leading to JAK1 and STAT2-mediated phosphorylation of STAT3 (PubMed: 16982608). In turn, STAT3 translocates to the nucleus where it drives expression of anti-inflammatory mediators (PubMed: 18025162). Targets antigen-presenting cells (APCs) such as macrophages and monocytes and inhibits their release of pro- inflammatory cytokines including granulocyte-macrophage colony-stimulating factor /GM-CSF, granulocyte colony-stimulating factor/G- CSF, IL-1 alpha, IL-1 beta, IL-6, IL-8 and TNF (PubMed: 11564774 , PubMed: 1940799 , PubMed: 7512027). Also interferes with antigen presentation by reducing the expression of MHC-class II and co- stimulatory molecules, thereby inhibiting their ability to induce T cell activation (PubMed: 8144879). In addition, controls the inflammatory response of macrophages by reprogramming essential

metabolic pathways including mTOR signaling (By similarity).


Cellular Location

Secreted.

Tissue Location

Produced by a variety of cell lines, including T- cells, macrophages, mast cells and other cell types

Images



Western blot analysis of IL10 expression in Ramos cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.