

xCT Antibody

Rabbit mAb

Catalog # AP91476

Product Information

Application	WB, IF, ICC, IP
Primary Accession	Q9UPY5
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	CCBR1; Cysteine/glutamate transporter; SLC7A11; solute carrier family 7; xCT;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	55423

Additional Information

Dilution	WB 1:500~1:2000 ICC/IF 1:50~1:150 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human xCT
Description	Sodium-independent, high-affinity exchange of anionic amino acids with high specificity for anionic form of cystine and glutamate.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	SLC7A11 (HGNC:11059)
Function	Heterodimer with SLC3A2, that functions as an antiporter by mediating the exchange of extracellular anionic L-cystine and intracellular L-glutamate across the cellular plasma membrane (PubMed: 11133847 , PubMed: 11417227 , PubMed: 14722095 , PubMed: 15151999 , PubMed: 34880232 , PubMed: 35245456 , PubMed: 35352032). Provides L-cystine for the maintenance of the redox balance between extracellular L- cystine and L-cysteine and for the maintenance of the intracellular levels of glutathione that is essential for cells protection from oxidative stress (By similarity). The transport is sodium-independent, electroneutral with a stoichiometry of 1:1, and is drove by the high intracellular concentration of L-glutamate and the intracellular reduction of L-cystine (PubMed: 11133847 , PubMed: 11417227). In addition, mediates the import of L-kynurenine leading to anti-ferroptotic signaling propagation required to maintain L-cystine and glutathione homeostasis (PubMed: 35245456). Moreover, mediates N-acetyl-L-cysteine uptake into the placenta leading to subsequently down-regulation of pathways associated with oxidative stress, inflammation and apoptosis (PubMed: 34120018). In vitro can also transport L-aspartate

(PubMed:[11417227](#)). May participate in astrocyte and meningeal cell proliferation during development and can provide neuroprotection by promoting glutathione synthesis and delivery from non-neuronal cells such as astrocytes and meningeal cells to immature neurons (By similarity). Controls the production of pheomelanin pigment directly (By similarity).


Cellular Location

Cell membrane; Multi-pass membrane protein. Cell projection, microvillus membrane; Multi-pass membrane protein. Note=Localized to the microvillous membrane of the placental syncytiotrophoblast.

Tissue Location

Expressed in term placenta and primary term cytotrophoblast (PubMed:34120018). Expressed mainly in the brain, but also in pancreas (PubMed:11417227).

Images



Western blot analysis of xCT expression in (1) HepG2 cell lysate; (2) Mouse brain lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.