

EAAT3 Antibody

Rabbit mAb

Catalog # AP91955

Product Information

Application	WB, IHC, IF, ICC, IHF
Primary Accession	P43005
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	SLC1A1, EAAC1, EAAT3, Eaac-1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	57100

Additional Information

Dilution	WB 1:1000~1:5000 IHC 1:100~1:500 ICC/IF 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human EAAT3
Description	Transports L-glutamate and also L- and D-aspartate. Essential for terminating the postsynaptic action of glutamate by rapidly removing released glutamate from the synaptic cleft. Acts as a symport by cotransporting sodium. Negatively regulated by ARL6IP5.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	SLC1A1 (HGNC:10939)
Function	Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate (PubMed: 21123949 , PubMed: 26690923 , PubMed: 33658209 , PubMed: 7521911 , PubMed: 7914198 , PubMed: 8857541). Can also transport L-cysteine (PubMed: 21123949). Functions as a symporter that transports one amino acid molecule together with two or three Na(+) ions and one proton, in parallel with the counter-transport of one K(+) ion (PubMed: 26690923 , PubMed: 33658209 , PubMed: 7521911 , PubMed: 8857541). Mediates Cl(-) flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na(+) symport (PubMed: 26690923 , PubMed: 8857541). Plays an important role in L- glutamate and L-aspartate reabsorption in renal tubuli (PubMed: 21123949). Plays a redundant role in the rapid removal of released glutamate from the synaptic cleft, which is essential for terminating the postsynaptic action of glutamate (By similarity). Contributes to glutathione biosynthesis and protection against oxidative

stress via its role in L-glutamate and L-cysteine transport (By similarity).
Negatively regulated by ARL6IP5 (By similarity).


Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P43003}. Apical cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P43003}. Synapse, synaptosome {ECO:0000250|UniProtKB:P51906}. Early endosome membrane {ECO:0000250|UniProtKB:P51906}. Late endosome membrane {ECO:0000250|UniProtKB:P51906}. Recycling endosome membrane {ECO:0000250|UniProtKB:P51906}

Tissue Location

Expressed in all tissues tested including liver, muscle, testis, ovary, retinoblastoma cell line, neurons and brain (in which there was dense expression in substantia nigra, red nucleus, hippocampus and in cerebral cortical layers)

Images



Western blot analysis of EAAT3 expression in Human fetal brain lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.