

IKZF3 Antibody

Rabbit mAb Catalog # AP92226

Product Information

Application WB, IHC, IF, FC, ICC, IHF

Primary Accession Q9UKT9

Reactivity Rat, Human, Mouse

Clonality Monoclonal

Other Names AIO; Aiolos; IKZF3; ZNFN1A3;

IsotypeRabbit IgGHostRabbitCalculated MW58023

Additional Information

Dilution WB 1:1000~1:5000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human IKZF3

Description Transcription factor that plays an important role in the regulation of

lymphocyte differentiation. Plays an essential role in regulation of B-cell differentiation, proliferation and maturation to an effector state. Involved in regulating BCL2 expression and controlling apoptosis in T-cells in an

IL2-dependent manner.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name IKZF3

Synonyms ZNFN1A3

Function Transcription factor that plays an important role in the regulation of

lymphocyte differentiation. Plays an essential role in regulation of B-cell differentiation, proliferation and maturation to an effector state. Involved in regulating BCL2 expression and controlling apoptosis in T-cells in an

Il 2 descendent manner


IL2-dependent manner.

Cellular Location Nucleus. Cytoplasm [Isoform 3]: Nucleus [Isoform 14]: Nucleus. Cytoplasm

Tissue Location Expressed most strongly in peripheral blood leukocytes, the spleen, and the

thymus.

Images



Western blot analysis of IKZF3 expression in Ramos cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.