

Aly Antibody

Rabbit mAb

Catalog # AP92228

Product Information

Application	WB, IHC, IF, FC, ICC, IP, IHF
Primary Accession	Q86V81
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	ALY; ALY/REF; BEF; REF; Tho4; thoc4;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	26888

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:40 FC 1:100
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Aly
Description	Component of the THO subcomplex of the TREX complex. The TREX complex specifically associates with spliced mRNA and not with unspliced pre-mRNA. It is recruited to spliced mRNAs by a transcription-independent mechanism. Binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	ALYREF
Synonyms	ALY, BEF, THOC4
Function	Functions as an mRNA export adapter; component of the transcription/export (TREX) complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced pre-mRNA (PubMed: 15833825 , PubMed: 15998806 , PubMed: 17190602). TREX is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm via the TAP/NXF1 pathway (PubMed: 15833825 , PubMed: 15998806 , PubMed: 17190602). Involved in the nuclear export of intronless mRNA; proposed to be recruited to intronless mRNA by ATP-bound

DDX39B (PubMed:[17984224](#)). Plays a key role in mRNP recognition and mRNA packaging by bridging the mRNP-bound EJC and the TREX core complex (PubMed:[37020021](#)). TREX recruitment occurs via an interaction between ALYREF/THOC4 and the cap-binding protein NCBP1 (PubMed:[15833825](#), PubMed:[15998806](#), PubMed:[17190602](#), PubMed:[37020021](#)). Required for TREX complex assembly and for linking DDX39B to the cap-binding complex (CBC) (PubMed:[15998806](#), PubMed:[17984224](#), PubMed:[37020021](#)). Binds mRNA which is thought to be transferred to the NXF1-NXT1 heterodimer for export (TAP/NXF1 pathway) (PubMed:[11675789](#), PubMed:[11707413](#), PubMed:[11979277](#), PubMed:[15833825](#), PubMed:[15998806](#), PubMed:[17190602](#), PubMed:[18364396](#), PubMed:[22144908](#), PubMed:[22893130](#), PubMed:[23222130](#), PubMed:[25662211](#)). In conjunction with THOC5 functions in NXF1-NXT1 mediated nuclear export of HSP70 mRNA; both proteins enhance the RNA binding activity of NXF1 and are required for NXF1 localization to the nuclear rim (PubMed:[19165146](#)). Involved in mRNA export of C5-methylcytosine (m5C)-containing mRNAs: specifically recognizes and binds m5C mRNAs and mediates their nucleo-cytoplasmic shuttling (PubMed:[28418038](#)). Acts as a chaperone and promotes the dimerization of transcription factors containing basic leucine zipper (bZIP) domains and thereby promotes transcriptional activation (PubMed:[10488337](#)). Involved in transcription elongation and genome stability (PubMed:[12438613](#)).


Cellular Location

Nucleus. Nucleus speckle Cytoplasm Note=Colocalizes with the core EJC, NXF1 and DDX39B in the nucleus and nuclear speckles. Travels to the cytoplasm as part of the exon junction complex (EJC) bound to mRNA (PubMed:[19324961](#)). Localizes to regions surrounding nuclear speckles known as perispeckles in which TREX complex assembly seems to occur (PubMed:[23826332](#))

Tissue Location

Expressed in a wide variety of cancer types.

Images



Western blot analysis of Aly expression in HEK293 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.