

ACHA4 Antibody

Rabbit mAb

Catalog # AP92399

Product Information

Application	WB, IP
Primary Accession	P43681
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	Neuronal acetylcholine receptor subunit alpha-4; ACH4; AChR; Acra4; BFNC; CHRNA4; EBN; EBN1; NACHR; NACRA4;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	69957

Additional Information

Dilution	WB 1:1000~1:5000 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human ACHA4
Description	After binding acetylcholine, the AChR responds by an extensive change in conformation that affects all subunits and leads to opening of an ion-conducting channel across the plasma membrane.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	CHRNA4 (HGNC:1958)
Synonyms	NACRA4
Function	Component of neuronal acetylcholine receptors (nAChRs) that function as pentameric, ligand-gated cation channels with high calcium permeability among other activities. nAChRs are excitatory neurotransmitter receptors formed by a collection of nAChR subunits known to mediate synaptic transmission in the nervous system and the neuromuscular junction. Each nAChR subunit confers differential attributes to channel properties, including activation, deactivation and desensitization kinetics, pH sensitivity, cation permeability, and binding to allosteric modulators (PubMed: 22361591 , PubMed: 27698419 , PubMed: 29720657 , PubMed: 38454578). CHRNA4 forms heteropentameric neuronal acetylcholine receptors with CHRNB2 and CHRNB4, as well as CHRNA5 and CHRNB3 as accessory subunits. Is the most abundant nAChR subtype expressed in the central nervous system (PubMed: 16835356 , PubMed: 22361591 , PubMed: 27698419 ,

PubMed:[29720657](#), PubMed:[38454578](#)). Found in two major stoichiometric forms, (CHRNA4)₃:(CHRN2)₂ and (CHRNA4)₂:(CHRN2)₃, the two stoichiometric forms differ in their unitary conductance, calcium permeability, ACh sensitivity and potentiation by divalent cation (PubMed:[27698419](#), PubMed:[29720657](#), PubMed:[38454578](#)). Involved in the modulation of calcium-dependent signaling pathways, influences the release of neurotransmitters, including dopamine, glutamate and GABA (By similarity).

Cellular Location

Synaptic cell membrane {ECO:0000250|UniProtKB:O70174}; Multi-pass membrane protein. Cell membrane {ECO:0000250|UniProtKB:O70174}; Multi-pass membrane protein

Images



Western blot analysis of ACHA4 expression in SH-SY5Y cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.