

5HT7 Receptor Antibody

Rabbit mAb Catalog # AP92505

Product Information

Application WB Primary Accession P34969

Reactivity Rat, Human, Mouse

Clonality Monoclonal

Other Names 5 HT 7; 5 HT X; 5-HT-7; 5-HT-X; 5-HT7; 5HT7; htr7;

IsotypeRabbit IgGHostRabbitCalculated MW53555

Additional Information

Dilution WB 1:500~1:2000

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human 5HT7 Receptor

Description This is one of the several different receptors for 5-hydroxytryptamine

(serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. The activity of this receptor is mediated by G

proteins that stimulate adenylate cyclase.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name HTR7 {ECO:0000303 | Ref.3, ECO:0000312 | HGNC:HGNC:5302}

Function G-protein coupled receptor for 5-hydroxytryptamine (serotonin), a biogenic

hormone that functions as a neurotransmitter, a hormone and a mitogen (PubMed:<u>35714614</u>, PubMed:<u>8226867</u>). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G

proteins) and modulates the activity of downstream effectors

(PubMed:35714614, PubMed:8226867). HTR7 is coupled to G(s) G alpha

proteins and mediates activation of adenylate cyclase activity


(PubMed:<u>35714614</u>).

Cellular Location Cell membrane; Multi-pass membrane protein

Tissue Location [Isoform A]: Predominant isoform in spleen, caudate and hippocampus.

[Isoform D]: Minor isoform in terms of expression.

Images



Western blot analysis of 5HT7 Receptor expression in U-87MG cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.