

TRAF2 Antibody

Rabbit mAb

Catalog # AP92633

Product Information

Application	WB, IHC, IF, FC, ICC, IP, IHF
Primary Accession	Q12933
Reactivity	Human
Clonality	Monoclonal
Other Names	TNF receptor associated factor 2; TRAF2; TRAP; TRAP3;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	55859

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human TRAF2
Description	Regulates activation of NF-kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	TRAF2 {ECO:0000303 PubMed:28489822, ECO:0000312 HGNC:HGNC:12032}
Function	E3 ubiquitin-protein ligase that regulates activation of NF- kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis (PubMed: 10346818 , PubMed: 11784851 , PubMed: 12917689 , PubMed: 15383523 , PubMed: 18981220 , PubMed: 19150425 , PubMed: 19810754 , PubMed: 19918265 , PubMed: 19937093 , PubMed: 20047764 , PubMed: 20064526 , PubMed: 20385093 , PubMed: 20577214 , PubMed: 22212761). Catalyzes 'Lys-63'-linked ubiquitination of target proteins, such as BIRC3, IKBKE, MLST8, RIPK1 and TICAM1 (PubMed: 23453969 , PubMed: 28489822). Is an essential constituent of several E3 ubiquitin- protein ligase complexes, where it promotes the ubiquitination of target proteins by bringing them into contact with other E3 ubiquitin ligases (PubMed: 15383523 , PubMed: 18981220). Regulates BIRC2 and BIRC3 protein levels by inhibiting their autoubiquitination and subsequent degradation; this does not depend on the TRAF2 RING-type zinc finger domain (PubMed: 11907583 , PubMed: 19506082). Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR (PubMed: 15121867). In complex with BIRC2 or BIRC3, promotes ubiquitination of IKBKE

(PubMed:[23453969](#)). Acts as a regulator of mTORC1 and mTORC2 assembly by mediating 'Lys-63'-linked ubiquitination of MLST8, thereby inhibiting formation of the mTORC2 complex, while facilitating assembly of the mTORC1 complex (PubMed:[28489822](#)). Required for normal antibody isotype switching from IgM to IgG (By similarity).

Cellular Location

Cytoplasm

Images



Western blot analysis of TRAF2 expression in Raji cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.