

# CCDC98 Antibody

Rabbit mAb Catalog # AP92807

### **Product Information**

**Application** WB, IF, FC, ICC, IP

Primary Accession

Reactivity

Clonality

Q6UWZ7

Human

Monoclonal

Other Names ABRA1; ABRAXAS; FAM175A;

IsotypeRabbit IgGHostRabbitCalculated MW46663

## **Additional Information**

**Dilution** WB 1:500~1:2000 ICC/IF 1:50~1:200 IP 1:50 FC 1:50

**Purification** Affinity-chromatography

**Immunogen** A synthesized peptide derived from human CCDC98

**Description** Component of the BRCA1-A complex, a complex that specifically recognizes

'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at

double-strand breaks (DSBs).

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

#### **Protein Information**

Name ABRAXAS1 ( HGNC:25829)

**Function** Involved in DNA damage response and double-strand break (DSB) repair.


Component of the BRCA1-A complex, acting as a central scaffold protein that assembles the various components of the complex and mediates the recruitment of BRCA1. The BRCA1-A complex specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesion sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at DSBs. This complex also possesses deubiquitinase activity that specifically

removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX.

**Cellular Location** Nucleus Note=Localizes at sites of DNA damage at double-strand breaks

(DSBs)

## **Images**



Western blot analysis of CCDC98 expression in MCF7 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.