

# LCE4A Rabbit Polyclonal Antibody

LCE4A Rabbit Polyclonal Antibody

Catalog # AP93377

## Product Information

|                          |                        |
|--------------------------|------------------------|
| <b>Application</b>       | IHC, IF                |
| <b>Primary Accession</b> | <a href="#">Q5TA78</a> |
| <b>Reactivity</b>        | Rat, Human, Mouse      |
| <b>Host</b>              | Polyclonal, Rabbit,IgG |
| <b>Clonality</b>         | Polyclonal             |
| <b>Calculated MW</b>     | 9980                   |

## Additional Information

|                           |  |
|---------------------------|--|
| <b>Gene ID</b>            | 199834   |
| <b>Other Names</b>        | Late cornified envelope protein 4A, Late envelope protein 8, Small proline-rich-like epidermal differentiation complex protein 4A, LCE4A, LEP8, SPRL4A |
| <b>Dilution</b>           | IHC~~1:100~500 IF~~1:50~200  |
| <b>Storage Conditions</b> | -20°C  |

## Protein Information

|                        |   |
|------------------------|---|
| <b>Name</b>            | LCE4A   |
| <b>Synonyms</b>        | LEP8, SPRL4A  |
| <b>Function</b>        | Precursors of the cornified envelope of the stratum corneum.  |
| <b>Tissue Location</b> | Skin-specific. Expression was readily detected in adult trunk skin, adult arm skin, fetal skin, penal skin, vulva, esophagus and tongue. Not expressed in the cervix, rectum, lung, colon, or placenta. |

## Images



Immunohistochemical analysis of paraffin-embedded human spleen tissue. 1,primary Antibody was diluted at 1:200(4° overnight). 2, Sodium citrate pH 6.0 was used for antigen retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.