

Androgen Receptor(AR-V7 specific) (6K19) Rabbit Monoclonal Antibody

Androgen Receptor(AR-V7 specific) (6K19) Rabbit Monoclonal Antibody
Catalog # AP93784

Product Information

Application	WB, IF, FC, ICC
Primary Accession	P10275-3
Reactivity	Human
Clonality	Monoclonal

Additional Information

Dilution	WB~~1:1000 IF~~1:50~200 FC~~1:10~50 ICC~~N/A
Storage Conditions	-20°C


Protein Information

Background

The androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. Expansion of the polyglutamine tract from the normal 9-34 repeats to the pathogenic 38-62 repeats causes spinal bulbar muscular atrophy (SBMA, also known as Kennedy's disease). Mutations in this gene are also associated with complete androgen insensitivity (CAIS). Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jan 2017]

Images

Western blot analysis of extracts from HepG2,HCT116 cells using AP93784 at 1:1000.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.