10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SCHIP1 (4X12) Mouse Monoclonal antibody

SCHIP1 (4X12) Mouse Monoclonal antibody
Catalog # AP93875

Product Information

Application	WB, IHC, IF
Primary Accession	P0DPB3
Reactivity	Human, Mouse
Clonality	Monoclonal
Calculated MW	53480

Additional Information

Gene ID	29970
Other Names	Schwannomin-interacting protein 1 {ECO:0000312|HGNC:HGNC:15678}, SCHIP-1, SCHIP1 (HGNC:15678)
Dilution	WB~~1:1000 IHC~~1:100~500 IF~~1:50~200
Storage Conditions	-20℃

Protein Information

Name	SCHIP1 (HGNC:15678)
Cellular Location	Cytoplasm.
Tissue Location	Preferentially expressed in brain, skeletal muscles and heart. Also expressed in detected in pancreas, kidney, liver, lung, and placenta.

Images



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-SCHIP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, AP93875)



Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-SCHIP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, AP93875)



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-SCHIP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, AP93875)



Anti-SCHIP1 mouse monoclonal antibody (AP93875) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY SCHIP1 .



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SCHIP1 (Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SCHIP1(Cat# AP93875). Positive lysates (100ug) and (20ug) can be purchased separately from biodragon.