

# UBE2Q1 Rabbit pAb

UBE2Q1 Rabbit pAb

Catalog # AP94048

## Product Information

---

|                     |   |
|---------------------|---|
| Primary Accession   | <a href="#">Q7Z7E8.1</a>  |
| Reactivity          | Human   |
| Host                | Rabbit  |
| Clonality           | Polyclonal  |
| Calculated MW       | 46 KDa  |
| Physical State      | Liquid  |
| Immunogen           | KLH conjugated synthetic peptide derived from human UBE2Q1  |
| Epitope Specificity | 321-422/422   |
| Isotype             | IgG   |
| Purity              | affinity purified by Protein A  |
| Buffer              | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.   |
| Important Note      | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. |

## Additional Information

---

|          |   |
|----------|---|
| Dilution | Flow-Cyt=2 $\mu$ g /Test  |
| Format   | 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce  |
| Storage  | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |

## Protein Information

---

## Background


---

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

## Images

---

cells:mouse splenocyte Concentration: 2  $\mu$ g/10<sup>6</sup> cells,  
4°C incubate 30min.



Blank control: Mouse spleen. Primary Antibody (green line): Rabbit Anti-UBE2Q1 antibody (AP94048) Dilution: 2  $\mu\text{g}$  / $10^6$  cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1  $\mu\text{g}$  /test. Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.