

phospho-Serine/Threonine Rabbit pAb

phospho-Serine/Threonine Rabbit pAb

Catalog # AP94054

Product Information

Application	IHC-P, IHC-F, IF, E
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	KLH conjugated synthesised phosphopeptide contain Phosphoserine and Phosphothreonine
Epitope Specificity	(p-S)(p-T)-NH ₂
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Protein phosphorylation provides a signalling system that can be thought of as a kind of protein on/off switch for many cellular signalling pathways. Phosphorylation is observed on serine, threonine, tyrosine and histidine residues. Cellular networks underlying phosphorylation can be very complex and often occurs on multiple distinct sites on a given protein. Phospho-specific antibodies are becoming critical reagents both for basic research and for clinical diagnosis.


Additional Information

Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

Protein phosphorylation provides a signalling system that can be thought of as a kind of protein on/off switch for many cellular signalling pathways. Phosphorylation is observed on serine, threonine, tyrosine and histidine residues. Cellular networks underlying phosphorylation can be very complex and often occurs on multiple distinct sites on a given protein. Phospho-specific antibodies are becoming critical reagents both for basic research and for clinical diagnosis.

Images



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phosphoserine,threonine) Polyclonal Antibody, Unconjugated (AP94054) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.