

Artemin Rabbit pAb

Artemin Rabbit pAb Catalog # AP94144

Product Information

Application WB, IHC-P, IHC-F, IF

Primary Accession Q9Z0L2 Reactivity Mouse Host Rabbit Clonality Polyclonal Calculated MW 23726 **Physical State** Liquid

Immunogen KLH conjugated synthetic peptide derived from mouse Artemin

141-224/224 **Epitope Specificity**

Isotype IgG

affinity purified by Protein A **Purity**

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION

SIMILARITY Belongs to the TGF-beta family. GDNF subfamily. **SUBUNIT** Homodimer; disulfide-linked. Binds to RET.

This product as supplied is intended for research use only, not for use in **Important Note**

human, therapeutic or diagnostic applications.

This gene encodes a secreted ligand of the glial cell line-derived neurotrophic **Background Descriptions**

factor (GDNF) subfamily and TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta

receptors leading to recruitment and activation of SMAD family transcription

factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. This protein signals through the RET receptor and GFR alpha 3 coreceptor, and supports the survival of a number of peripheral neuron populations and at least one population of dopaminergic CNS neurons. This protein has also been shown to promote tumor growth, metastasis, and drug

resistance in mammary carcinoma. [provided by RefSeq, Aug 2016]

Additional Information

11876 Gene ID

Other Names Artemin, Artn {ECO:0000303 | PubMed:9883723,

ECO:0000312 | MGI:MGI:1333791}

Cochlea. Expressed at higher level in sesorineural epithelium than in the Target/Specificity

modiolus region or substantia nigra.

Dilution WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500 Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name Artn {ECO:0000303|PubMed:9883723, ECO:0000312|MGI:MGI:1333791}

Function Growth factor that supports the survival of sensory and sympathetic


peripheral neurons in culture and also supports the survival of dopaminergic neurons of the ventral mid-brain (By similarity). Acts by binding to its coreceptor, GFRA3, leading to autophosphorylation and activation of the RET receptor (PubMed:15204970). Strong attractant of gut hematopoietic cells thus promoting the formation Peyer's patch-like structures, a major component of the gut-associated lymphoid tissue (PubMed:17322904).

Cellular Location Secreted.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images




Sample: Testis (Mouse) Lysate at 40 ug Primary: Anti-Artemin (AP94144) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 12/19 kD Observed band size: 19 kD



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Artemin) Polyclonal Antibody, Unconjugated (AP94144) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody



incubation with (Artemin) Polyclonal Antibody, Unconjugated (AP94144) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.