

# phospho-AMPK alpha 1 (Ser356) Rabbit pAb

phospho-AMPK alpha 1 (Ser356) Rabbit pAb

Catalog # AP94172

## Product Information

---

<b>Application</b>	WB, IHC-P, IHC-F, IF
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	64 KDa
<b>Physical State</b>	Liquid
<b>Immunogen</b>	KLH conjugated Synthesised phosphopeptide derived from human AMPK alpha 1 around the phosphorylation site of Ser356
<b>Epitope Specificity</b>	AT(p-S)PP
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
<b>SUBCELLULAR LOCATION</b>	Cytoplasm. Nucleus. Note=In response to stress, recruited by p53/TP53 to specific promoters.
<b>SIMILARITY</b>	Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. SNF1 subfamily. Contains 1 protein kinase domain.
<b>SUBUNIT</b>	AMPK is a heterotrimer of an alpha catalytic subunit (PRKAA1 or PRKAA2), a beta (PRKAB1 or PRKAB2) and a gamma non-catalytic subunits (PRKAG1, PRKAG2 or PRKAG3). Interacts with FNIP1 and FNIP2.
<b>Post-translational modifications</b>	Ubiquitinated. Phosphorylated at Thr-183 by STK11/LKB1 in complex with STE20-related adapter-alpha (STRADA) pseudo kinase and CAB39. Also phosphorylated at Thr-183 by CAMKK2; triggered by a rise in intracellular calcium ions, without detectable changes in the AMP/ATP ratio. CAMKK1 can also phosphorylate Thr-183, but at much lower level. Dephosphorylated by protein phosphatase 2A and 2C (PP2A and PP2C). Phosphorylated by ULK1 and ULK2; leading to negatively regulate AMPK activity and suggesting the existence of a regulatory feedback loop between ULK1, ULK2 and AMPK.
<b>DISEASE</b>	Defects in CRYAB are the cause of myofibrillar alpha-B crystallin-related (MFM-CRYAB) [MIM:608810]. A neuromuscular disorder that results in weakness of the proximal and distal limb muscles, weakness of the neck, velopharynx and trunk muscles, hypertrophic cardiomyopathy, and cataract in a subset of patients.
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>Background Descriptions</b>	The protein encoded by this gene belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio. AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008]


## Additional Information

<b>Target/Specificity</b>	Lens as well as other tissues.
<b>Dilution</b>	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=1 ug /Test
<b>Format</b>	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


## Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.


## Images



Paraformaldehyde-fixed, paraffin embedded (Human cervical carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-AMPK alpha 1 (Ser356)) Polyclonal Antibody, Unconjugated (AP94172) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (Mouse liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-AMPK alpha 1 (Ser356)) Polyclonal Antibody, Unconjugated (AP94172) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-AMPK alpha 1 (Ser356)) Polyclonal Antibody, Unconjugated (AP94172) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.


Sample: Hela(Human) CellLysate at 30 ug MCF-7(Human) CellLysate at 30 ug Primary: Anti-phospho-AMPK alpha 1 (Ser356) (AP94172) at 1/500 dilution Secondary:



IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 63 kD Observed band size: 63 kD



Sample: NIH/3T3(Mouse) Cell Lysate at 30 ug Primary:  
Anti-phospho-AMPK alpha 1 (Ser356) (AP94172) at 1/500  
dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at  
1/20000 dilution Predicted band size: 63 kD Observed  
band size: 63 kD



Blank control(black line):HepG2. Primary Antibody (green  
line): Rabbit Anti-phospho-AMPK alpha 1 (Ser356)  
antibody (AP94172) Dilution:1ug/Test; Secondary  
Antibody(white blue line): Goat anti-rabbit IgG-AF488  
Dilution: 0.5ug/Test. Isotype control(orange line): Normal  
Rabbit IgG Protocol The cells were fixed with 4% PFA  
(10min at room temperature)and then permeabilized  
with 90% ice-cold methanol for 20 min at -20°C, The cells  
were then incubated in 5%BSA to block non-specific  
protein-protein interactions for 30 min at room  
temperature .Cells stained with Primary Antibody for 30  
min at room temperature. The secondary antibody used  
for 40 min at room temperature. Acquisition of 20,000  
events was performed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.