

Phospho-RPS6KA1 (Ser380) Rabbit pAb

Phospho-RPS6KA1 (Ser380) Rabbit pAb Catalog # AP94191

Product Information

ApplicationWBReactivityHumanHostRabbitClonalityPolyclonalCalculated MW81 KDaPhysical StateLiquid

Immunogen KLH conjugated Synthesised phosphopeptide derived from human RPS6KA1

around the phosphorylation site of Ser380

Epitope Specificity GF(p-S)FV **Isotype** IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nucleus. Cytoplasm.

SIMILARITY Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family.

S6 kinase subfamily.Contains 1 AGC-kinase C-terminal domain.Contains 2

protein kinase domains.

SUBUNIT Forms a complex with either MAPK1/ERK2 or MAPK3/ERK1 in quiescent cells.

Transiently dissociates following mitogenic stimulation. Interacts with

ETV1/ER81 and FGFR1.

Post-translational Activated by phosphorylation at Ser-221 by PDPK1. Autophosphorylated on Ser-380, as part of the activation process. May be phosphorylated at Thr-35!

Ser-380, as part of the activation process. May be phosphorylated at Thr-359 and Ser-363 by MAPK1/ERK2 and MAPK3/ERK1.N-terminal myristoylation results in an activated kinase in the absence of added growth factors.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions Rsk1 is a member of a family of 90kDa ribosomal protein S6 kinases, which

includes Rsk1, Rsk2 and Rsk3. These are broadly expressed serine/threonine

protein kinases activated in response to mitogenic stimuli, including

extracellular signal regulated protein kinases Erk1 and Erk2. Rsk1 is activated by MAPK in vitro and in vivo via phosphorylation. Active Rsks appear to play a major role in transcriptional regulation by translocating to the nucleus and

phosphorylating c Fos and CREB.

Additional Information

Dilution WB=1:500-2000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce


Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample: A431(Human) Cell Lysate at 30 ug Primary: Anti-Phospho-RPS6KA1(Ser380) (AP94191) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 81 kD Observed band size: 81 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.