

PCID2 Rabbit pAb

PCID2 Rabbit pAb
Catalog # AP94205

Product Information

Application	WB
Primary Accession	Q5JVF3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46030
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human PCID2
Epitope Specificity	101-200/399
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the CSN12 family. Contains 1 PCI domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA _i and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. May favor virus entry in case of infection with herpes simplex virus 1 (HSV1) or herpes simplex virus 2 (HSV2).

Additional Information

Gene ID	55795
Other Names	PCI domain-containing protein 2, CSN12-like protein, PCID2
Dilution	WB=1:500-2000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	PCID2
Function	Required for B-cell survival through the regulation of the expression of cell-cycle checkpoint MAD2L1 protein during B cell differentiation (By similarity). As a component of the TREX-2 complex, involved in the export of mRNAs to the cytoplasm through the nuclear pores (PubMed: 22307388). Binds and stabilizes BRCA2 and is thus involved in the control of R-loop-associated DNA damage and transcription-associated genomic instability (PubMed: 24896180). Blocks the activity of the SRCAP chromatin remodeling complex by interacting with SRCAP complex member ZNHIT1 and inhibiting its interaction with the complex (By similarity). This prevents the deposition of histone variant H2AZ1/H2A.Z at the nucleosomes of key lymphoid fate regulator genes which suppresses their expression and restricts lymphoid lineage commitment (By similarity).
Cellular Location	Cytoplasm. Nucleus, nuclear pore complex

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample: Lane 1: Rat HeLa tissue lysates Lane 2: Rat Placenta tissue lysates Primary: Anti-PCID2 (AP94205) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 46 kDa Observed band size: 45 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.