

UPF1 Recombinant Mouse mAb

UPF1 Recombinant Mouse mAb

Catalog # AP94322

Product Information

Application	WB, IF, ICC
Host	Rabbit
Clonality	Recombinant
Physical State	Liquid
Isotype	IgG1, Kappa
Purity	affinity purified by Protein G
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Cytoplasm, P-body. Note=Hyperphosphorylated form is targeted to the P-body, while unphosphorylated protein is distributed throughout the cytoplasm.
SIMILARITY	Belongs to the DNA2/NAM7 helicase family. Contains 1 C2H2-type zinc finger.
SUBUNIT	Found in a post-splicing messenger ribonucleoprotein (mRNP) complex. Associates with the exon junction complex (EJC). Associates with the SGM1C complex; is phosphorylated by the complex kinase component SGM1. Interacts with UPF2, UPF3A and UPF3B. Interacts with EST1A and SLBP. Interacts (when hyperphosphorylated) with PNRC2. Interacts with EIF2C1, EIF2C2 and GSPT2.
Post-translational modifications	Phosphorylated by SMG1; required for formation of mRNA surveillance complexes. Phosphorylated upon DNA damage, probably by ATM or ATR.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	In eukaryotes, it is essential to have the ability to detect and degrade transcripts that lack full coding potential. Nonsense-mediated RNA decay (NMD) protects the organism by avoiding the translation of truncated peptides with dominant negative or deleterious gain-of-function potential. Rent1, a mammalian ortholog of Upflp, is essential for embryonic viability (1-3). Rent1 (also designated regulator of nonsense transcripts and HUpf1) contains an N-terminal zinc finger-like domain, NTPase domains and a region comprised of domains that define Rent1 as a superfamily group I helicase.


Additional Information

Target/Specificity	Ubiquitous.
Dilution	WB=1:500-1:1000, ICC/IF=1:50-1:200
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Cell line: HeLa Fixative: 4% Paraformaldehyde
Permeabilization: 0.1% TritonX-100 Primary ab dilution:
1:200 Primary incubation condition: 4°C overnight
Secondary ab: Goat Anti-Mouse IgG Nuclear counter
stain: DAPI (Blue) Comment: Color green is the positive
signal for AP94322



Blocking buffer: 5% NFDM/TBST Primary ab dilution:
1:1000 Primary ab incubation condition: 4°C overnight
Secondary ab: Goat Anti-Mouse IgG H&L (HRP) Lysate:
Jurkat, Raji, SH-SY5Y, 293 Protein loading quantity: 20 µg
Exposure time: 30 s Predicted MW: 124 kDa Observed
MW: 130 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.