

GNB3 Recombinant Mouse mAb

GNB3 Recombinant Mouse mAb Catalog # AP94520

Product Information

Application WB, IHC-P, IHC-F, IF, ICC

Host Rabbit
Clonality Recombinant
Physical State Liquid
Isotype IgG1, Kappa

Purity affinity purified by Protein G

Buffer0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. **SIMILARITY**Belongs to the WD repeat G protein beta family. Contains 7 WD repeats. **SUBUNIT**G proteins are composed of 3 units, alpha, beta and gamma. Interacts with

RASD2.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions G proteins are composed of 3 units, alpha, beta and gamma and are involved

in the modulation and transduction of various transmembrane signalling systems. There are families of related genes for each subunit. GNB3 encodes

a beta subunit. Beta subunits are required for GTPase activity, for replacement of GDP by GTP, regulate alpha subunits and for G

protein-effector interactions. A single-nucleotide polymorphism (C825T) in this gene is associated with essential hypertension and obesity and is also associated with the occurrence of the splice variant GNB3-s, which appears to

have increased activity.

Additional Information

Dilution WB=1:500-1:1000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:50,IF=0

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When


reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Tissue: Human liver cancer Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Mouse)(sp-0024) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for AP94520



Cell line: HepG2 Fixative: 4% Paraformaldehyde Permeabilization: 0.1% TritonX-100 Primary ab dilution: 1:50 Primary incubation condition: 4°C overnight Secondary ab: Goat Anti-Rabbit IgG Nuclear counter stain: DAPI (Blue) Comment: Color green is the positive signal for AP94520

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.