

# LHB Recombinant Rabbit mAb

LHB Recombinant Rabbit mAb

Catalog # AP94638

## Product Information

---

<b>Application</b>	WB, IHC-P, IHC-F, IF, ICC
<b>Host</b>	Rabbit
<b>Clonality</b>	Recombinant
<b>Calculated MW</b>	16 KDa
<b>Physical State</b>	Liquid
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
<b>SUBCELLULAR LOCATION</b>	Secreted.
<b>SIMILARITY</b>	Belongs to the glycoprotein hormones subunit beta family.
<b>SUBUNIT</b>	Heterodimer of a common alpha chain and a unique beta chain which confers biological specificity to thyrotropin, lutropin, follitropin and gonadotropin.
<b>DISEASE</b>	Hypogonadism LHB-related (HGON-LHB) [MIM:152780]: Characterized by infertility and pseudohermaphroditism. Note=The disease is caused by mutations affecting the gene represented in this entry.
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>Background Descriptions</b>	Luteinizing Hormone is a member of the glycoprotein hormone family. Glycoprotein hormones are heterodimers consisting of a common alpha subunit and an unique beta subunit which confers biological specificity. LH is expressed in the pituitary gland and promotes spermatogenesis and ovulation by stimulating the testes and ovaries to synthesize steroids. Mutations in the Luteinizing Hormone (LHB) gene are associated with hypogonadism which is characterized by infertility and pseudohermaphroditism.

## Additional Information

---


<b>Target/Specificity</b>	Pituitary gland.
<b>Dilution</b>	WB=1:500-1:2000,IHC-P=1:100-500,IHC-F=,ICC/IF=1:50-1:100,IF=0,Flow-Cyt=1:50-1:100
<b>Format</b>	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Background


---

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.


## Images




Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Rabbit IgG H&L (HRP) Lysate: 1: Rat pituitary, 2: Mouse pituitary Protein loading quantity: 20 µg Exposure time: 10 s Predicted MW: 15 kDa Observed MW: 15 kDa



Cell line: HeLa Fixative: 4% Paraformaldehyde Permeabilization: 0.1% TritonX-100 Primary ab dilution: 1:50 Primary incubation condition: 4°C overnight Secondary ab: Goat Anti-Rabbit IgG Nuclear counter stain: DAPI (Blue) Comment: Color green is the positive signal for AP94638




Cell line: PANC-1 Fixative: 100% Ice-cold methanol Permeabilization: 0.1% TritonX-100 Primary ab dilution: 1:50 Primary incubation condition: 4°C overnight Secondary ab: Goat Anti-Rabbit IgG Nuclear counter stain: DAPI (Blue) Comment: Color green is the positive signal for AP94638



Cell line: HeLa Fixative: 4% Paraformaldehyde Permeabilization: 90% Methanol Primary ab dilution: 1:100 Secondary ab: Goat anti Rabbit IgG Unlabelled control: The cell without incubation with primary antibody and secondary antibody (Black line). Isotype control: Rabbit monoclonal IgG (Blue line). Comment: Line red is the positive signal for AP94638

Tissue: Human pituitary Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High



temperature and high pressure Retrieval buffer:  
Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:1000  
Primary ab incubation condition: 1 hour at room  
temperature Secondary ab: SP Kit(Rabbit) (sp-0023)  
Counter stain: Hematoxylin (Blue) Comment: Color brown  
is the positive signal for AP94638

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.