

# DRD2 Rabbit pAb

DRD2 Rabbit pAb Catalog # AP94806

#### **Product Information**

ApplicationWBReactivityHumanHostRabbitClonalityPolyclonalCalculated MW51 KDaPhysical StateLiquid

Immunogen KLH conjugated synthetic peptide derived from human DRD2

Epitope Specificity 109-170/444

**Isotype** IgG

**Purity** affinity purified by Protein A

**Buffer** 0.01M TBS(pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

**SUBCELLULAR LOCATION** Cell membrane.

**SIMILARITY** Belongs to the G-protein coupled receptor 1 family.

DISEASE

Defects in DRD2 are associated with dystonia type 11 (DYT11) [MIM:159900]; also known as alcohol-responsive dystonia. DYT11 is a myoclonic dystonia.

Dystonia is defined by the presence of sustained involuntary muscle contractions, often leading to abnormal postures. DYT11 is characterized by involuntary lightning jerks and dystonic movements and postures alleviated by alcohol. Inheritance is autosomal dominant. The age of onset, pattern of

body involvement, presence of myoclonus and response to alcohol are all

variable.

**Important Note** This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

**Background Descriptions** This gene encodes the D2 subtype of the dopamine receptor. This G-protein

coupled receptor inhibits adenylyl cyclase activity. A missense mutation in this gene causes myoclonus dystonia; other mutations have been associated with schizophrenia. Alternative splicing of this gene results in two transcript variants encoding different isoforms. A third variant has been described, but it has not been determined whether this form is normal or due to aberrant

splicing. [provided by RefSeq, Jul 2008]

### **Additional Information**

**Dilution** WB=1:500-2000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When


reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

## **Background**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

## **Images**



Sample: Lane 1: Mouse Cerebrum tissue lysates Lane 2: Rat Cerebrum tissue lysates Primary: Anti-DRD2 (AP94806) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 51 kDa Observed band size: 50 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.