

# LDHA Rabbit pAb

LDHA Rabbit pAb Catalog # AP94839

#### **Product Information**

**Application** WB, IHC-P, IHC-F, IF

Primary Accession P00338

**Reactivity** Rat, Human, Mouse

Host Rabbit
Clonality Polyclonal
Calculated MW 36689
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived from human LDHA

**Isotype** IgG

**Purity** affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

**SUBCELLULAR LOCATION** Cytoplasm.

**SIMILARITY** Belongs to the LDH/MDH superfamily. LDH family.

**SUBUNIT** Homotetramer. **Post-translational** ISGylated.

modifications

**DISEASE** Glycogen storage disease 11 (GSD11) [MIM:612933]: A metabolic disorder that

results in exertional myoglobinuria, pain, cramps and easy fatigue. Note=The disease is caused by mutations affecting the gene represented in this entry. This product as supplied is intended for research use only, not for use in

Important Note This product as supplied is intended for research use only, not

human, therapeutic or diagnostic applications.

#### **Additional Information**

**Gene ID** 3939

Other Names L-lactate dehydrogenase A chain, LDH-A, 1.1.1.27, Cell proliferation-inducing

gene 19 protein, LDH muscle subunit, LDH-M, Renal carcinoma antigen

NY-REN-59, LDHA (HGNC:6535)

**Target/Specificity** Metabolism of carbohydrates, Dehydrogenase

Kits, NADP/NADPH, Carbohydrate metabolism, Energy

Metabolism, Cancer, Energy Metabolism

**Dilution** WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:400-800,IF=1:10-50

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

#### **Protein Information**

Name LDHA ( HGNC:6535)

**Function** Interconverts simultaneously and stereospecifically pyruvate and lactate

with concomitant interconversion of NADH and NAD(+).

Cellular Location Cytoplasm.


**Tissue Location** Predominantly expressed in anaerobic tissues such as skeletal muscle and

liver.

### **Background**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.


## **Images**




25 ug total protein per lane of various lysates (see on figure) probed with LDHA polyclonal antibody, unconjugated (AP94839) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Paraformaldehyde-fixed, paraffin embedded Mouse Skeletal muscle; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with LDHA Polyclonal Antibody, Unconjugated (AP94839) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Human Skeletal muscle; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with LDHA Polyclonal Antibody, Unconjugated (AP94839) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Mouse Heart; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with LDHA Polyclonal Antibody, Unconjugated (AP94839) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.





Paraformaldehyde-fixed, paraffin embedded Human Lung Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with LDHA Polyclonal Antibody, Unconjugated (AP94839) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.

Paraformaldehyde-fixed, paraffin embedded Human Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with LDHA Polyclonal Antibody, Unconjugated (AP94839) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.