

EFHA2 Rabbit pAb

EFHA2 Rabbit pAb Catalog # AP94841

Product Information

Application IHC-P, IHC-F, IF

Primary Accession Q86XE3

Reactivity Rat, Rabbit, Dog, Horse

Host Rabbit
Clonality Polyclonal
Calculated MW 60711
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived from human EFHA2

Epitope Specificity 451-530/530

Isotype IgG

Purity affinity purified by Protein A

Buffer0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. **SUBCELLULAR LOCATION**Mitochondrion (By similarity). Membrane; Single-pass membrane protein

(Potential).

SIMILARITY Contains 2 EF-hand domains.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Additional Information

Gene ID 286097

Other Names Calcium uptake protein 3, mitochondrial, hMICU3, EF-hand

domain-containing family member A2, MICU3

{ECO:0000303|PubMed:30699349, ECO:0000312|HGNC:HGNC:27820}

Dilution IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name MICU3 {ECO:0000303 | PubMed:30699349,

ECO:0000312 | HGNC:HGNC:27820}

Function Tissue-specific calcium sensor of the mitochondrial calcium uniporter (MCU)

channel, which specifically regulates MCU channel activity in the central

nervous system and skeletal muscle (PubMed:<u>29725115</u>). Senses calcium level

via its EF-hand domains: compared to MICU1 and MICU2, MICU3 has a higher affinity for calcium (PubMed:29725115). MICU1 and MICU3 form a disulfide-linked heterodimer that stimulates and inhibits MCU activity, depending on the concentration of calcium (PubMed:29725115). At low calcium levels, MICU1 occludes the pore of the MCU channel, preventing mitochondrial calcium uptake (PubMed:29725115). At higher calcium levels, calcium- binding to MICU1 and MICU3 induces a conformational change that weakens MCU-MICU1 interactions and moves the MICU1-MICU3 heterodimer away from the pore, allowing calcium permeation through the MCU channel (PubMed: 29725115). The high calcium affinity of MICU3 lowers the calcium threshold necessary for calcium permeation through the MCU channel (PubMed: <u>29725115</u>). The MICU1-MICU3 heterodimer promotes flexibility of neurotransmission in neuronal cells by enhancing mitochondrial calcium uptake in presynapses (PubMed:29725115). It is also required to increase mitochondrial calcium uptake in skeletal muscle cells, thereby increasing ATP production (By similarity).

Cellular Location

Mitochondrion intermembrane space {ECO:0000250|UniProtKB:Q8IYU8}. Mitochondrion inner membrane {ECO:0000250|UniProtKB:Q8IYU8}. Note=Recruited to the mitochondrial inner membrane via its association with the uniplex complex {ECO:0000250|UniProtKB:Q8IYU8}

Tissue Location

Specifically expressed in the central nervous system and skeletal muscle.

Background

MICU3 (Mitochondrial Calcium Uptake Family Member 3) is a Protein Coding gene. Gene Ontology (GO) annotations related to this gene include calcium ion binding. An important paralog of this gene is MICU2.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.