

# GALNT5 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9976a

## Product Information

---

<b>Application</b>	IHC-P, FC, WB, E
<b>Primary Accession</b>	<a href="#">Q7Z7M9</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB24611
<b>Calculated MW</b>	106266
<b>Antigen Region</b>	24-53

## Additional Information

---

<b>Gene ID</b>	11227
<b>Other Names</b>	Polypeptide N-acetylgalactosaminyltransferase 5, Polypeptide GalNAc transferase 5, GalNAc-T5, pp-GaNTase 5, Protein-UDP acetylgalactosaminyltransferase 5, UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase 5, GALNT5
<b>Target/Specificity</b>	This GALNT5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 24-53 amino acids from the N-terminal region of human GALNT5.
<b>Dilution</b>	IHC-P~~1:100~500 FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	GALNT5 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	GALNT5
<b>Function</b>	Catalyzes the initial reaction in O-linked oligosaccharide biosynthesis, the

transfer of an N-acetyl-D-galactosamine residue to a serine or threonine residue on the protein receptor. Has activity toward EA2 peptide substrate, but has a weak activity toward Muc2 or Muc1b substrates (By similarity).

## Cellular Location

Golgi apparatus membrane; Single-pass type II membrane protein


## Background

GALNT5 can catalyze the initial reaction in O-linked oligosaccharide biosynthesis, the transfer of an N-acetyl-D-galactosamine residue to a serine or threonine residue on the protein receptor. GALNT5 has activity toward EA2 peptide substrate, but it has a weak activity toward Muc2 or Muc1b substrates.


## References

Simmons, A.D., et al. Hum. Mol. Genet. 8(12):2155-2164(1999)


## Images



Western blot analysis of lysates from A431, K562 cell line (from left to right), using GALNT5 Antibody (N-term)(Cat. #AP9976a). AP9976a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



GALNT5 Antibody (N-term) (Cat. #AP9976a) IHC analysis in formalin fixed and paraffin embedded human normal lung tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the GALNT5 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



GALNT5 Antibody (N-term) (Cat. #AP9976a) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

## Citations

---

- [The Glycoprotein Mucin-1 Negatively Regulates GalNAc Transferase 5 Expression in Pancreatic Cancer.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.