

# Anti-TfR Reference Antibody (Jr-141)

Recombinant Antibody  
Catalog # APR10179

## Product Information

---

|                          |                            |
|--------------------------|----------------------------|
| <b>Application</b>       | FC, Kinetics, Animal Model |
| <b>Primary Accession</b> | <a href="#">P02786</a>     |
| <b>Reactivity</b>        | Human, Mouse               |
| <b>Clonality</b>         | Monoclonal                 |
| <b>Isotype</b>           | IgG1                       |
| <b>Calculated MW</b>     | 84871                      |

## Additional Information

---

|                                  |  |
|----------------------------------|--|
| <b>Target/Specificity</b>        | TfR  |
| <b>Endotoxin<br/>Conjugation</b> | Unconjugated   |
| <b>Expression system</b>         | CHO Cell   |
| <b>Format</b>                    | Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column. |

## Protein Information


---

|                 |   |
|-----------------|---|
| <b>Name</b>     | TFRC  |
| <b>Function</b> | Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes (PubMed: <a href="#">26214738</a> ). Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the hereditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C- terminal binding site. Positively regulates T and B cell proliferation through iron uptake (PubMed: <a href="#">26642240</a> ). Acts as a lipid sensor that regulates mitochondrial fusion by regulating activation of the JNK pathway (PubMed: <a href="#">26214738</a> ). When dietary levels of stearate (C18:0) are low, promotes activation of the JNK pathway, resulting in HUWE1- mediated ubiquitination and subsequent degradation of the mitofusin MFN2 and inhibition of mitochondrial fusion (PubMed: <a href="#">26214738</a> ). When dietary levels of stearate (C18:0) are high, TFRC stearylolation inhibits activation of the JNK pathway and thus degradation of the mitofusin MFN2 (PubMed: <a href="#">26214738</a> ). Mediates uptake of NICOL1 into fibroblasts where it may regulate extracellular matrix production (By similarity). |


Cellular Location

Cell membrane; Single-pass type II membrane protein Melanosome.  
Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV


Images



Anti-TfR Reference Antibody (Jr-141) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-TfR Reference Antibody (Jr-141) is more than 95% ,determined by SEC-HPLC.



Immobilized human Transferrin R , His Tag at 2 µg/mL can bind Anti-TfR Reference Antibody (Jr-141),EC50=0.0093 µg/mL

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.