



Anti-ERBB1 / EGFR / HER1 Reference Antibody (Duke D2C7)

Recombinant Antibody Catalog # APR10895

Product Information

Application FC, Kinetics, Animal Model

Primary Accession
Reactivity
Human
Clonality
Monoclonal
Isotype
IgG1
Calculated MW
134277

Additional Information

Target/Specificity ERBB1 / EGFR / HER1

Endotoxin

Conjugation Unconjugated

Expression system CHO Cell

Format Purified monoclonal antibody supplied in PBS, pH6.0, without

preservative. This antibody is purified through a protein A column.

Protein Information

Name EGFR (HGNC:3236)

Synonyms ERBB, ERBB1, HER1

Function Receptor tyrosine kinase binding ligands of the EGF family and activating

several signaling cascades to convert extracellular cues into appropriate cellular responses (PubMed:<u>10805725</u>, PubMed:<u>27153536</u>, PubMed:<u>2790960</u>, PubMed:<u>35538033</u>). Known ligands include EGF, TGFA/TGF- alpha, AREG, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding

EGF (PubMed: 12297049, PubMed: 15611079, PubMed: 17909029,

PubMed:20837704, PubMed:27153536, PubMed:2790960, PubMed:7679104, PubMed:8144591, PubMed:9419975). Ligand binding triggers receptor homoand/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules (PubMed:27153536). May also activate the NF-kappa-B signaling cascade (PubMed:11116146). Also directly phosphorylates other proteins like RGS16, activating its GTPase

activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling (PubMed:11602604). Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin (PubMed:11483589). Positively regulates cell migration via interaction with CCDC88A/GIV which retains EGFR at the cell membrane following ligand stimulation, promoting EGFR signaling which triggers cell migration (PubMed:20462955). Plays a role in enhancing learning and memory performance (By similarity). Plays a role in mammalian pain signaling (long-lasting hypersensitivity) (By similarity).


Cellular Location

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein Golgi apparatus membrane; Single-pass type I membrane protein. Nucleus membrane; Single-pass type I membrane protein. Endosome. Endosome membrane. Nucleus. Note=In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER (PubMed:17909029, PubMed:20674546). Endocytosed upon activation by ligand (PubMed:17182860, PubMed:17909029, PubMed:27153536, PubMed:2790960). Colocalized with GPER1 in the nucleus of estrogen agonist-induced cancer-associated fibroblasts (CAF) (PubMed:20551055)


Tissue Location

Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.

Images



Anti-ERBB1 / EGFR / HER1 Reference Antibody (Duke D2C7) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-ERBB1 / EGFR / HER1 Reference Antibody (Duke D2C7)is more than 95% ,determined by SEC-HPLC.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.