

# Anti-Mouse IgG (gamma 1, 2a, 2b and 3 chain) (Rhodamine Conjugated) Secondary Antibody

Rabbit Polyclonal, Rhodamine (TRITC) Catalog # ASR2739

### **Product Information**

**Description** Anti-MOUSE IgG (gamma 1, 2a, 2b and 3 chain) (RABBIT) Antibody Rhodamine

Conjugated

**Host** Rabbit

**Conjugate** Rhodamine (TRITC)

**FP Value** 3.1 moles Rhodamine (TRITC) per mole of IgG

Target SpeciesMouseClonalityPolyclonalPhysical StateLyophilized

Host Isotype IgG

Target Isotype IgG (gamma 1, 2a, 2b and 3 chain)

**Buffer** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Immunogen** highly purified mouse IgG gamma 1, gamma 2a, gamma 2b and gamma 3

proteins

**Reconstitution Volume** 1.0 mL

**Reconstitution Buffer** Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

**Preservative** 0.01% (w/v) Sodium Azide

#### **Additional Information**

Shipping Condition Ambient

**Application Note** Anti-Mouse IgG subclass is also suitable for multiplex analysis, including

multicolor imaging, utilizing various commercial platforms.

Purity Anti-Mouse IgG subclass pan reactive Secondary Antibody was prepared

from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. This product shows balanced reactivity to Mouse IgG1, IgG2a, IgG2b and IgG3 proteins and is suitable to screen IgG class hybridoma clones. Minimal cross reactivity is observed against other

Mouse immunoglobulin classes or light chain proteins.

**Storage Condition** Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.


**Precautions Note**This product is for research use only and is not intended for therapeutic or

diagnostic applications.

# **Background**

Rhodamine Conjugated Secondary Antibodies are ideal for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting and are available in a variety of formats and conjugate types. When choosing a secondary antibody, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

## **Images**



Indirect ELISA of Rabbit Anti-Mouse IgG (gamma 1, 2a, 2b, and 3) antibody. Antigen: purified mouse IgG heavy and light chains. Coating amount: 0.1 µg per well. Primary antibody: Rabbit Anti-Mouse IgG (Gamma 1, 2a, 2b, and 3) HRP conjugated Antibody. Dilution series: 2-fold. Substrate: TMB (p/n TMBE-0100).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.