

CUTL1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant CUTL1.

Catalog # AT1684a

Product Information

Application	WB, IHC, IF, IP, E
Primary Accession	Q13948
Other Accession	NM_001913
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG1 kappa
Clone Names	2A10
Calculated MW	77455

Additional Information


Gene ID	1523
Other Names	Protein CASP, CUX1, CUTL1
Target/Specificity	CUTL1 (NP_001904.2, 521 a.a. ~ 620 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IHC~~1:100~500 IF~~1:50~200 IP~~N/A E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	CUTL1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

Background


The protein encoded by this gene is a member of the homeodomain family of DNA binding proteins. It may regulate gene expression, morphogenesis, and differentiation and it may also play a role in the cell cycle progression. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some of these variants has not been determined.

References

1. Neocortical Layer Formation of Human Developing Brains and Lissencephalies: Consideration of Layer-Specific Marker Expression. Saito T, Hanai S, Takashima S, Nakagawa E, Okazaki S, Inoue T, Miyata R, Hoshino K, Akashi T, Sasaki M, Goto YI, Hayashi M, Itoh M. Cereb Cortex. 2010 Jul 12. [Epub ahead of print]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .




Western Blot analysis of CUX1 expression in transfected 293T cell line by CUTL1 monoclonal antibody (M01), clone 2A10.

Lane 1: CUX1 transfected lysate(77.2 KDa).

Lane 2: Non-transfected lysate.





Immunoperoxidase of monoclonal antibody to CUTL1 on formalin-fixed paraffin-embedded human malignant lymphoma, diffuse large B tissue. [antibody concentration 5 ug/ml]



Immunofluorescence of monoclonal antibody to CUTL1 on HeLa cell. [antibody concentration 10 ug/ml]

Immunoprecipitation of CUX1 transfected lysate using anti-CUX1 monoclonal antibody and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with CUX1 MaxPab rabbit polyclonal antibody.



Detection limit for recombinant GST tagged CUTL1 is approximately 0.03ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.