

ICA1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant ICA1. Catalog # AT2467a

Product Information

Application WB, E
Primary Accession Q05084
Other Accession NM_022307
Reactivity Human
Host mouse
Clonality monoclonal
Isotype IgG2a Kappa
Clone Names 6G11

Clone Names 6G11 Calculated MW 54645

Additional Information

Gene ID 3382

Other Names Islet cell autoantigen 1, 69 kDa islet cell autoantigen, ICA69, Islet cell

autoantigen p69, ICAp69, p69, ICA1

Target/Specificity ICA1 (NP_071682, 1 a.a. ~ 110 a.a) partial recombinant protein with GST tag.

MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions ICA1 Antibody (monoclonal) (M01) is for research use only and not for use in

diagnostic or therapeutic procedures.

Background


This gene encodes a protein with an arfaptin homology domain that is found both in the cytosol and as membrane-bound form on the Golgi complex and immature secretory granules. This protein is believed to be an autoantigen in insulin-dependent diabetes mellitus and primary Sjogren's syndrome. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized.

References


Common variants in FOXP1 are associated with generalized vitiligo. Jin Y, et al. Nat Genet, 2010 Jul. PMID

20526340. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. ICA69 is a novel Rab2 effector regulating ER-Golgi trafficking in insulinoma cells. Buffa L, et al. Eur J Cell Biol, 2008 Apr. PMID 18187231. Large-scale mapping of human protein-protein interactions by mass spectrometry. Ewing RM, et al. Mol Syst Biol, 2007. PMID 17353931. The LIFEdb database in 2006. Mehrle A, et al. Nucleic Acids Res, 2006 Jan 1. PMID 16381901.

Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 KDa) .



Western Blot analysis of ICA1 expression in transfected 293T cell line by ICA1 monoclonal antibody (M01), clone 6G11.

Lane 1: ICA1 transfected lysate (Predicted MW: 54.6 KDa). Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged ICA1 is approximately 0.3ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.