

IL6ST Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant IL6ST. Catalog # AT2523a

Product Information

ApplicationWB, EPrimary AccessionP40189Other AccessionNM_002184ReactivityHumanHostmouseClonalitymonoclonalIsotypeIgG2a Kappa

Clone Names 4A4

Additional Information

Other Names Interleukin-6 receptor subunit beta, IL-6 receptor subunit beta, IL-6R subunit

beta, IL-6R-beta, IL-6RB, CDw130, Interleukin-6 signal transducer, Membrane glycoprotein 130, gp130, Oncostatin-M receptor subunit alpha, CD130, IL6ST

Target/Specificity IL6ST (NP_002175, 23 a.a. ~ 122 a.a) partial recombinant protein with GST tag.

MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions IL6ST Antibody (monoclonal) (M02) is for research use only and not for use in

diagnostic or therapeutic procedures.

Background


The protein encoded by this gene is a signal transducer shared by many cytokines, including interleukin 6 (IL6), ciliary neurotrophic factor (CNTF), leukemia inhibitory factor (LIF), and oncostatin M (OSM). This protein functions as a part of the cytokine receptor complex. The activation of this protein is dependent upon the binding of cytokines to their receptors. vIL6, a protein related to IL6 and encoded by the Kaposi sarcoma-associated herpesvirus, can bypass the interleukin 6 receptor (IL6R) and directly activate this protein. Knockout studies in mice suggest that this gene plays a critical role in regulating myocyte apoptosis. Alternatively spliced transcript variants encoding distinct isoforms have been described. A related pseudogene has been identified on chromosome 17.

References

Evaluation of candidate stromal epithelial cross-talk genes identifies association between risk of serous

ovarian cancer and TERT, a cancer susceptibility hot-spot. Johnatty SE, et al. PLoS Genet, 2010 Jul 8. PMID 20628624. Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Continuous glycoprotein-130-mediated signal transducer and activator of transcription-3 activation promotes inflammation, left ventricular rupture, and adverse outcome in subacute myocardial infarction. Hilfiker-Kleiner D, et al. Circulation, 2010 Jul 13. PMID 20585009. Sortilin facilitates signaling of ciliary neurotrophic factor and related helical type 1 cytokines targeting the gp130/leukemia inhibitory factor receptor beta heterodimer. Larsen JV, et al. Mol Cell Biol, 2010 Sep. PMID 20584990. Crystal structure of the entire ectodomain of gp130: insights into the molecular assembly of the tall cytokine receptor complexes. Xu Y, et al. J Biol Chem, 2010 Jul 9. PMID 20489211.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.