

NR4A3 Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant NR4A3.

Catalog # AT3110a

Product Information

Application	WB, IF, E
Primary Accession	Q92570
Other Accession	NM_006981
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG2a Kappa
Clone Names	1E10
Calculated MW	68230

Additional Information

Gene ID	8013
Other Names	Nuclear receptor subfamily 4 group A member 3, Mitogen-induced nuclear orphan receptor, Neuron-derived orphan receptor 1, Nuclear hormone receptor NOR-1, NR4A3, CHN, CSMF, MINOR, NOR1, TEC
Target/Specificity	NR4A3 (NP_008912, 414 a.a. ~ 521 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IF~~1:50~200 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	NR4A3 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


Background

This gene encodes a member of the steroid-thyroid hormone-retinoid receptor superfamily. The encoded protein may act as a transcriptional activator. The protein can efficiently bind the NGFI-B Response Element (NBRE). Three different versions of extraskeletal myxoid chondrosarcomas (EMCs) are the result of reciprocal translocations between this gene and other genes. The translocation breakpoints are associated with Nuclear Receptor Subfamily 4, Group A, Member 3 (on chromosome 9) and either Ewing Sarcoma Breakpoint Region 1 (on chromosome 22), RNA Polymerase II, TATA Box-Binding Protein-Associated Factor, 68-KD (on chromosome 17), or Transcription factor 12 (on chromosome 15). Multiple transcript variants encoding different isoforms have been found for this gene.


References

Replicated association of the NR4A3 gene with smoking behavior in schizophrenia and in bipolar disorder. Novak G, et al. *Genes Brain Behav*, 2010 Jul 24. PMID 20659174. Deficiency of the NR4A orphan nuclear receptor NOR1 decreases monocyte adhesion and atherosclerosis. Zhao Y, et al. *Circ Res*, 2010 Aug 20. PMID 20558821. Fluorescence in situ hybridization analysis of extraskeletal myxoid chondrosarcomas using EWSR1 and NR4A3 probes. Noguchi H, et al. *Hum Pathol*, 2010 Mar. PMID 19775727. Preparation of polyclonal antibody specific for NOR1 and detection of its expression pattern in human tissues and nasopharyngeal carcinoma. Xiang B, et al. *Acta Biochim Biophys Sin (Shanghai)*, 2009 Sep. PMID 19727524. The hypoxia-inducible factor 1/NOR-1 axis regulates the survival response of endothelial cells to hypoxia. Martorell L, et al. *Mol Cell Biol*, 2009 Nov. PMID 19720740.


Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.62 KDa) .




NR4A3 monoclonal antibody (M02), clone 1E9 Western Blot analysis of NR4A3 expression in HeLa S3 NE ((Cat # AT3110a)



Immunofluorescence of monoclonal antibody to NR4A3 on HeLa cell. [antibody concentration 10 ug/ml]

Detection limit for recombinant GST tagged NR4A3 is approximately 0.3ng/ml as a capture antibody.



Citations

- [AP2-NR4A3 transgenic mice display reduced serum epinephrine because of increased catecholamine catabolism in adipose tissue.](#)
- [DNA-dependent protein kinase \(DNA-PK\) permits vascular smooth muscle cell proliferation through phosphorylation of the orphan nuclear receptor NOR1.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.