

HIST1H2BM Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5002

Product Information

Application	WB
Primary Accession	Q99879
Reactivity	Human, Mouse
Predicted	Mouse
Host	Rabbit
Clonality	polyclonal
Calculated MW	13989
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	8342
Antigen Region	10-44
Other Names	Histone H2B type 1-M, Histone H2Be, H2B/e, HIST1H2BM, H2BFE
Dilution	WB~1:1000
Target/Specificity	This HIST1H2BM antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 10-44 amino acids from the N-terminal region of human HIST1H2BM.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	HIST1H2BM Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	H2BC14 (HGNC:4750)
Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription

regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location

Nucleus. Chromosome.


Background

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

References

Albig W.,et al.Hum. Genet. 101:284-294(1997).
Marzluff W.F.,et al.Genomics 80:487-498(2002).
Mungall A.J.,et al.Nature 425:805-811(2003).
Lubec G.,et al.Submitted (MAR-2007) to UniProtKB.
Cheung W.L.,et al.Cell 113:507-517(2003).

Images



All lanes : Anti-HIST1H2BM Antibody (N-term) at 1:2000 dilution
Lane 1: PC-3 whole cell lysate
Lane 2: HepG2 whole cell lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 14 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.