

UBE2D2 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5140

Product Information

Application	WB
Primary Accession	P62837
Reactivity	Human, Rat
Predicted	Mouse
Host	Rabbit
Clonality	polyclonal
Calculated MW	16735
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	7322
Antigen Region	44-76
Other Names	Ubiquitin-conjugating enzyme E2 D2, Ubiquitin carrier protein D2, Ubiquitin-conjugating enzyme E2(17)KB 2, Ubiquitin-conjugating enzyme E2-17 kDa 2, Ubiquitin-protein ligase D2, p53-regulated ubiquitin-conjugating enzyme 1, UBE2D2, PUBC1, UBC4, UBC5B, UBCH4, UBCH5B
Dilution	WB~~1:500
Target/Specificity	This UBE2D2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 44-76 amino acids from the N-terminal region of human UBE2D2.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	UBE2D2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	UBE2D2
-------------	--------

Synonyms

PUBC1, UBC4, UBC5B, UBCH4, UBCH5B

Function

Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins (PubMed:[10329681](#), PubMed:[18042044](#), PubMed:[18703417](#), PubMed:[20061386](#), PubMed:[20403326](#), PubMed:[20525694](#), PubMed:[26475854](#), PubMed:[28322253](#)). Catalyzes 'Lys-48'-linked polyubiquitination (PubMed:[10329681](#), PubMed:[18042044](#), PubMed:[18359941](#), PubMed:[18703417](#), PubMed:[20061386](#), PubMed:[20403326](#), PubMed:[20525694](#), PubMed:[26475854](#)). Mediates the selective degradation of short-lived and abnormal proteins (PubMed:[10329681](#), PubMed:[18042044](#), PubMed:[18359941](#), PubMed:[18703417](#), PubMed:[20061386](#), PubMed:[20403326](#), PubMed:[20525694](#), PubMed:[26475854](#)). Functions in the E6/E6-AP-induced ubiquitination of p53/TP53 (PubMed:[15280377](#)). Mediates ubiquitination of PEX5 and SQSTM1 and autoubiquitination of STUB1 and TRAF6 (PubMed:[18359941](#), PubMed:[28322253](#)). Involved in the signal-induced conjugation and subsequent degradation of NFKBIA, FBXW2-mediated GCM1 ubiquitination and degradation, MDM2-dependent degradation of p53/TP53 and the activation of MAVS in the mitochondria by RIGI in response to viral infection (PubMed:[18703417](#), PubMed:[20403326](#)). Essential for viral activation of IRF3 (PubMed:[19854139](#)).


Background

Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins. In vitro catalyzes 'Lys-48'-linked polyubiquitination. Mediates the selective degradation of short-lived and abnormal proteins. Functions in the E6/E6-AP-induced ubiquitination of p53/TP53. Mediates ubiquitination of PEX5 and autoubiquitination of STUB1 and TRAF6. Involved in the signal-induced conjugation and subsequent degradation of NFKBIA, FBXW2-mediated GCM1 ubiquitination and degradation, MDM2-dependent degradation of p53/TP53 and the activation of MAVS in the mitochondria by DDX58/RIG-I in response to viral infection. Essential for viral activation of IRF3.

References

Jensen J.P.,et al.J. Biol. Chem. 270:30408-30414(1995).
Rolfe M.,et al.Proc. Natl. Acad. Sci. U.S.A. 92:3264-3268(1995).
Guinn B.-A.,et al.Biochem. Biophys. Res. Commun. 335:1293-1304(2005).
Yin Y.,et al.Submitted (OCT-2000) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



All lanes : Anti-UBE2D2 Antibody (N-term) at 1:2000 dilution Lane 1: F9 whole cell lysate Lane 2: NIH/3T3 whole cell lysate Lane 3: SW480 whole cell lysate Lane 4: 293 whole cell lysate Lane 5: Jurkat whole cell lysate Lane 6: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 17 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.