

PDIA6 Antibody (Center K159)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5167

Product Information

Application	IF, IHC-P, FC, WB
Primary Accession	Q15084
Reactivity	Human, Rat
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Antigen Region	144-172
Other Names	PDIA6; ERP5; P5; TXNDC7; Protein disulfide-isomerase A6; Endoplasmic reticulum protein 5; Protein disulfide isomerase P5; Thioredoxin domain-containing protein 7
Dilution	IF~~1:100 IHC-P~~1:100 FC~~1:10~50 WB~~1:1000
Target/Specificity	This PDIA6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 144-172 amino acids from the Central region of human PDIA6.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PDIA6 Antibody (Center K159) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Background


Protein disulfide isomerases (EC 5.3.4.1), such as PDIA6, are endoplasmic reticulum (ER) resident proteins that catalyze formation, reduction, and isomerization of disulfide bonds in proteins and are thought to play

a role in folding of disulfide-bonded proteins.


References

Hayano,T.,Gene 164 (2), 377-378 (1995)


Images




Western blot analysis of lysates from HT-1080, K562 cell line, human liver tissue, HepG2, C6 cell line (from left to right), using PDIA6 Antibody (Center K159) (Cat. #AW5167). AW5167 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody.




Formalin-fixed and paraffin-embedded human brain tissue reacted with PDIA6 Antibody (Center K159), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.




Flow cytometric analysis of HeLa cells using PDIA6 Antibody (Center K159) (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Immunohistochemical analysis of paraffin-embedded H. liver section using PDIA6 Antibody (Center K159) (Cat#AW5167). AW5167 was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded H. brain section using PDIA6 Antibody (Center K159)(Cat#AW5167). AW5167 was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Fluorescent image of HepG2 cells stained with XAF1 PDIA6 Antibody (Center K159)(Cat#AW5167). AW5167 was diluted at 1:100 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody (green). DAPI was used to stain the cell nuclear (blue). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.