

HMGCS1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AW5269

Product Information

Application FC, IHC-P, WB **Primary Accession** Q01581

Other Accession <u>P17425, Q8|ZK9, P13704, P23228</u>

Reactivity Mouse, Rat, Human **Predicted** Hamster, Chicken

Host Rabbit
Clonality Polyclonal
Calculated MW 57294
Isotype Rabbit IgG
Antigen Source HUMAN

Additional Information

Gene ID 3157

Antigen Region 290-317

Other Names HMGCS1; HMGCS; Hydroxymethylglutaryl-CoA synthase, cytoplasmic;

3-hydroxy-3-methylglutaryl coenzyme A synthase

Dilution FC~~1:10~50 IHC-P~~1:100~500 WB~~ 1:1000

Target/Specificity This HMGCS1 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 290-317 amino acids from the Central

region of human HMGCS1.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation

followed by dialysis against PBS.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions HMGCS1 Antibody (Center) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name HMGCS1 (HGNC:5007)

Synonyms HMGCS

Function

Catalyzes the condensation of acetyl-CoA with acetoacetyl-CoA to form HMG-CoA, which is converted by HMG-CoA reductase (HMGCR) into mevalonate, a precursor for cholesterol synthesis.

Cellular Location

Cytoplasm.


Background

HMGCS1 condenses acetyl-CoA with acetoacetyl-CoA to form HMG-CoA, which is the substrate for HMG-CoA reductase.


References

Vock, C., Cell. Physiol. Biochem. 22 (5-6), 515-524 (2008)


Images



Western blot analysis of lysates from mouse liver,rat liver,human liver tissue (from left to right), using HMGCS1 Antibody (Center)(Cat. #AW5269). AW5269 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.



Formalin-fixed and paraffin-embedded human lung carcinoma reacted with HMGCS1 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of hela cells using HMGCS1 Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.