

C1GALT1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AW5281

Product Information

Application WB **Primary Accession Q9NS00** Reactivity Human **Predicted** Rat Host Rabbit Clonality Polyclonal **Calculated MW** 42203 Isotype Rabbit IgG **Antigen Source** Human

Additional Information

Gene ID 56913

Antigen Region 115-144

Other Names Glycoprotein-N-acetylgalactosamine 3-beta-galactosyltransferase

1; Glycoprotein-N-acetylgalactosamine 3-beta-galactosyltransferase 1;

Glycoprotein-N-acetylgalactosamine 3-beta-galactosyltransferase 1; B3Gal-T8; Glycoprotein-N-acetylgalactosamine 3-beta-galactosyltransferase 1; Core 1

O-glycan T-synthase; Glycoprotein-N-acetylgalactosamine

3-beta-galactosyltransferase 1; Core 1

UDP-galactose:N-acetylgalactosamine-alpha-R beta 1, 3-galactosyltransferase 1; Glycoprotein-N-acetylgalactosamine 3-beta-galactosyltransferase 1; Core 1

beta1, 3-galactosyltransferase 1

Dilution WB~~1:1000

Target/Specificity This C1GALT1 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 115-144 amino acids from the Central

region of human C1GALT1.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions C1GALT1 Antibody (Center) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name C1GALT1

Function Glycosyltransferase that generates the core 1 O-glycan Gal-

beta1-3GalNAc-alpha1-Ser/Thr (T antigen), which is a precursor for many extended O-glycans in glycoproteins (PubMed:<u>11677243</u>). Plays a central role

in many processes, such as angiogenesis, thrombopoiesis and kidney

homeostasis development (By similarity).

Cellular Location Membrane {ECO:0000250 | UniProtKB:Q9JJ05}; Single- pass type II membrane

protein

Tissue Location Widely expressed. Highly expressed in kidney, heart, placenta and liver.


Background

Glycosyltransferase that generates the core 1 O-glycan Gal-beta1-3GalNAc-alpha1-Ser/Thr (T antigen), which is a precursor for many extended O-glycans in glycoproteins. Plays a central role in many processes, such as angiogenesis, thrombopoiesis and kidney homeostasis development.

References

Ju T., et al. J. Biol. Chem. 277:178-186(2002). Jensen M.P.A., et al. Submitted (JUN-1999) to the EMBL/GenBank/DDBJ databases. Hillier L.W., et al. Nature 424:157-164(2003). Scherer S.W., et al. Science 300:767-772(2003). Ju T., et al. Proc. Natl. Acad. Sci. U.S.A. 99:16613-16618(2002).

Images



Western blot analysis of lysates from mouse kidney and human kidney tissue (from left to right), using C1GALT1 Antibody (Center)(Cat. #AW5281). AW5281 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.