

WDR5 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5416

Product Information

Application	WB
Primary Accession	P61964
Other Accession	Q9V3J8 , Q498M4 , P61965 , Q2KIG2
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Antigen Region	321-354
Other Names	WD repeat-containing protein 5, BMP2-induced 3-kb gene protein, WDR5, BIG3
Dilution	WB~1:1000
Target/Specificity	This WDR5 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 321-354 amino acids from the C-terminal region of human WDR5.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	WDR5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Background


Contributes to histone modification. May position the N- terminus of histone H3 for efficient trimethylation at –Lys-4–. As part of the MLL1/MLL complex it is involved in methylation and dimethylation at

'Lys-4' of histone H3. H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation. As part of the NSL complex it may be involved in acetylation of nucleosomal histone H4 on several lysine residues. May regulate osteoblasts differentiation.

References

Young J.M., et al. Submitted (SEP-1998) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).
Wysocka J., et al. Genes Dev. 17:896-911(2003).
Hughes C.M., et al. Mol. Cell 13:587-597(2004).
Yokoyama A., et al. Mol. Cell. Biol. 24:5639-5649(2004).

Images



All lanes : Anti-WDR5 Antibody (C-term) at 1:1000 dilution
Lane 1: HeLa whole cell lysates Lane 2: NIH/3T3 whole cell lysates
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution
Predicted band size : 37 kDa
Blocking/Dilution buffer: 5% NFD/MTBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.