

SIRT3 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5443

Product Information

Application	WB
Primary Accession	Q9NTG7
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43573
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	23410
Antigen Region	250-279
Other Names	NAD-dependent protein deacetylase sirtuin-3, mitochondrial, hSIRT3, 351-, Regulatory protein SIR2 homolog 3, SIR2-like protein 3, SIRT3, SIR2L3
Dilution	WB~~1:1000
Target/Specificity	This SIRT3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 250-279 amino acids from the C-terminal region of human SIRT3.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SIRT3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SIRT3 {ECO:0000303 PubMed:12186850, ECO:0000312 HGNC:HGNC:14931}
Function	NAD-dependent protein deacetylase (PubMed: 12186850 , PubMed: 12374852 , PubMed: 16788062 , PubMed: 18680753 , PubMed: 18794531 , PubMed: 19535340 , PubMed: 23283301 , PubMed: 24121500 ,

PubMed:[24252090](#)). Activates or deactivates mitochondrial target proteins by deacetylating key lysine residues (PubMed:[12186850](#), PubMed:[12374852](#), PubMed:[16788062](#), PubMed:[18680753](#), PubMed:[18794531](#), PubMed:[23283301](#), PubMed:[24121500](#), PubMed:[24252090](#), PubMed:[38146092](#)). Known targets include ACSS1, IDH, GDH, SOD2, PDHA1, LCAD, SDHA, MRPL12 and the ATP synthase subunit ATP5PO (PubMed:[16788062](#), PubMed:[18680753](#), PubMed:[19535340](#), PubMed:[24121500](#), PubMed:[24252090](#), PubMed:[38146092](#)). Contributes to the regulation of the cellular energy metabolism (PubMed:[24252090](#)). Important for regulating tissue-specific ATP levels (PubMed:[18794531](#)). In response to metabolic stress, deacetylates transcription factor FOXO3 and recruits FOXO3 and mitochondrial RNA polymerase POLRMT to mtDNA to promote mtDNA transcription (PubMed:[23283301](#)). Acts as a regulator of ceramide metabolism by mediating deacetylation of ceramide synthases CERS1, CERS2 and CERS6, thereby increasing their activity and promoting mitochondrial ceramide accumulation (By similarity). Regulates hepatic lipogenesis (By similarity). Uses NAD(+) substrate imported by SLC25A47, triggering downstream activation of PRKAA1/AMPK- alpha signaling cascade that ultimately downregulates sterol regulatory element-binding protein (SREBP) transcriptional activities and ATP- consuming lipogenesis to restore cellular energy balance (By similarity). In addition to protein deacetylase activity, also acts as a protein-lysine deacylase by recognizing other acyl groups, such as benzoyl and lactoyl, leading to protein debenzoylation and delactylation, respectively (PubMed:[39524354](#), PubMed:[36896611](#), PubMed:[37720100](#)). Catalyzes debenzoylation of PPIF and ACLY (PubMed:[37720100](#)). Mediates delactylation of CCNE2 and 'Lys-16' of histone H4 (H4K16la) (PubMed:[36896611](#), PubMed:[37720100](#)).

Cellular Location

Mitochondrion matrix

Tissue Location

Widely expressed.

Background


SIRT3 is a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The SIRT3 is included in class I of the sirtuin family.

References

Hirschey, M.D., et al. Nature 464(7285):121-125(2010) Pillai, V.B., et al. J. Biol. Chem. 285(5):3133-3144(2010)
Kim, H.S., et al. Cancer Cell 17(1):41-52(2010)

Images

All lanes : Anti-SIRT3 Antibody (C-term) at 1:1000 dilution
Lane 1: HepG2 whole cell lysates Lane 2: Y79 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 44 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.