

EXOC3 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5482

Product Information

Application	WB, IHC-P
Primary Accession	O60645
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	87 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Antigen Region	524-557
Other Names	Exocyst complex component 3, Exocyst complex component Sec6, EXOC3, SEC6, SEC6L1
Dilution	WB~~1:1000 IHC-P~~1:100~500
Target/Specificity	This EXOC3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 524-557 amino acids from the C-terminal region of human EXOC3.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	EXOC3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Background

Component of the exocyst complex involved in the docking of exocytic vesicles with fusion sites on the plasma membrane.

References

Li J., et al. Mol. Cell. Proteomics 9:2517-2528(2010).


Jikuya H., et al. DNA Res. 10:49-57(2003).

Schmutz J., et al. Nature 431:268-274(2004).


Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Yu W., et al. Submitted (MAR-1998) to the EMBL/GenBank/DDBJ databases.

Images



AW5482 staining EXOC3 in Human spleen tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



All lanes : Anti-EXOC3 Antibody (C-term) at 1:1000 dilution
Lane 1: A431 whole cell lysates Lane 2: human brain lysates Lane 3: HeLa whole cell lysates Lane 4: U-2 OS whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 87 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.