

Mouse Gpi Antibody(C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5542

Product Information

Application	WB
Primary Accession	P06745
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	62767
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	14751
Antigen Region	532-558
Other Names	Glucose-6-phosphate isomerase, GPI, Autocrine motility factor, AMF, Neuroleukin, NLK, Phosphoglucose isomerase, PGI, Phosphohexose isomerase, PHI, Gpi, Gpi1
Dilution	WB~~1:1000
Target/Specificity	This Mouse Gpi antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 532-558 amino acids from the C-terminal region of mouse Gpi.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Mouse Gpi Antibody(C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Gpi {ECO:0000303 PubMed:7545951}
Function	In the cytoplasm, catalyzes the conversion of glucose-6- phosphate to fructose-6-phosphate, the second step in glycolysis, and the reverse reaction

during gluconeogenesis (PubMed:[2344351](#), PubMed:[7277315](#), PubMed:[8417789](#)). Besides its role as a glycolytic enzyme, also acts as a secreted cytokine: acts as an angiogenic factor (AMF) that stimulates endothelial cell motility (By similarity). Acts as a neurotrophic factor, neuroleukin, for spinal and sensory neurons (PubMed:[3352745](#), PubMed:[3764429](#)). It is secreted by lectin-stimulated T-cells and induces immunoglobulin secretion (PubMed:[3352745](#)).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P06744}. Secreted {ECO:0000250|UniProtKB:P06744}


Background

Besides its role as a glycolytic enzyme, mammalian GPI can function as a tumor-secreted cytokine and an angiogenic factor (AMF) that stimulates endothelial cell motility. GPI is also a neurotrophic factor (Neuroleukin) for spinal and sensory neurons.

References

Benveniste, P., et al. Cell Stem Cell 6(1):48-58(2010)
Repiso, A., et al. Anat Histol Embryol 37(5):380-382(2008)
Szuber, N., et al. Exp. Hematol. 36(7):773-785(2008)
Chandele, A., et al. J. Immunol. 180(8):5309-5319(2008)
Cuda, C.M., et al. J. Immunol. 179(11):7439-7447(2007)

Images



All lanes : Anti-Gpi Antibody (C-term) at 1:1000 dilution
Lane 1: mouse heart lysate Lane 2: mouse kidney lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 63 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.