

# Hsp60 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5684

## Product Information

---

<b>Application</b>	WB, IHC
<b>Primary Accession</b>	<a href="#">P10809</a>
<b>Other Accession</b>	<a href="#">P31081</a> , <a href="#">Q5ZL72</a> , <a href="#">P18687</a> , <a href="#">Q39727</a> , <a href="#">P63038</a> , <a href="#">Q5NVM5</a> , <a href="#">P63039</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Predicted</b>	Human, Mouse, Chicken, Dog
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	61 KDa
<b>Isotype</b>	Rabbit IgG
<b>Antigen Source</b>	HUMAN

## Additional Information

---

<b>Antigen Region</b>	396-430
<b>Other Names</b>	60 kDa heat shock protein, mitochondrial, 60 kDa chaperonin, Chaperonin 60, CPN60, Heat shock protein 60, HSP-60, Hsp60, HuCHA60, Mitochondrial matrix protein P1, P60 lymphocyte protein, HSPD1, HSP60
<b>Dilution</b>	WB~~1:40000 IHC~~1:1000
<b>Target/Specificity</b>	This Hsp60 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 396-430 amino acids from human Hsp60.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Hsp60 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

### Background

---


Implicated in mitochondrial protein import and macromolecular assembly. May facilitate the correct

folding of imported proteins. May also prevent misfolding and promote the refolding and proper assembly of unfolded polypeptides generated under stress conditions in the mitochondrial matrix.


## References

Jindal S., et al. *Mol. Cell. Biol.* 9:2279-2283(1989).  
Venner T.J., et al. *DNA Cell Biol.* 9:545-552(1990).  
Hansen J.J., et al. *Hum. Genet.* 112:71-77(2003).  
Tan J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.  
Ota T., et al. *Nat. Genet.* 36:40-45(2004).

## Images



All lanes : Anti-Hsp60 Antibody at 1:40000 dilution Lane 1: HeLa whole cell lysate Lane 2: A431 whole cell lysate Lane 3: A549 whole cell lysate Lane 4: HepG2 whole cell lysate Lane 5: NIH/3T3 whole cell lysate Lane 6: C2C12 whole cell lysate Lane 7: PC-12 whole cell lysate Lane 8: C6 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 60 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded Human liver section using Pink1 (Cat#AW5684). AW5684 was diluted at 1:1000 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.