

HCC-4/CCL16

Catalog # PVGS1126

Product Information

Primary Accession Species	O15467 Human
Sequence	Gln24-Gln120
Purity	> 97% as analyzed by SDS-PAGE > 97% as analyzed by HPLC
Endotoxin Level	
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human monocytes is in a concentration range of 10.0-100.0 ng/ml.
Expression System	E. coli
Theoretical Molecular Weight	11.2 kDa
Formulation	Lyophilized from a 0.2 μ m filtered solution in 20 mM PB, pH 7.4, 150 mM NaCl.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Additional Information

Other Names	C-C motif chemokine 16, Chemokine CC-4, HCC-4, Chemokine LEC, IL-10-inducible chemokine, LCC-1, Liver-expressed chemokine, Lymphocyte and monocyte chemoattractant, LMC, Monotactin-1, MTN-1, NCC-4, Small-inducible cytokine A16, CCL16, ILINCK, NCC4, SCYA16
Target Background	Human HCC-4, also named NCC-4, liver-expressed chemokine (LEC), and lymphocyte and monocyte chemoattractant (LMC), is a novel CC chemokine identified through bioinformatics. HCC-4 cDNA encodes a 120 amino acid (aa) residue precursor protein with a 23 aa residue predicted signal peptide that is cleaved to generate a 97 aa residue mature protein. HCC-4 is distantly related to other CC chemokines, exhibiting less than 30% sequence identity. Among these CC chemokines, HCC-4 has the most similarity to HCC-1. Two potential polyadenylation signals are present on the human HCC-4 gene, and as a result, two transcripts containing approximately 1,500 base pairs and 500

base pairs have been detected. HCC-4 is expressed weakly by some lymphocytes, including NK cells, T cells, and some T cell clones. The expression of HCC-4 in monocytes is highly upregulated in the presence of IL-10.

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.