

FGF-18

Catalog # PVGS1260

Product Information

Primary Accession 089101
Species Mouse

Sequence Glu28-Arg199

Purity > 95% as analyzed by SDS-PAGE

> 95% as analyzed by HPLC

Endotoxin Level

Expression System E. coli

Formulation Lyophilized after extensive dialysis against PBS.

Reconstitution It is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in

ddH₂O up to 100 □g/ml.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at lower than

-70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw

cycles.

Additional Information

Gene ID 14172

Other Names Fibroblast growth factor 18, FGF-18, zFGF5, Fgf18

Target Background Fibroblast Growth Factor 18 (FGF-18) is a pleiotropic cytokine belonging to the

heparin-binding FGF family, which has 23 different members. Structurally, FGF-18 is closely related to FGF-8 and FGF-17. Like other FGFs, FGF-18 can bind to different FGF receptors in vivo. FGF-18 is expressed in various tissues and has multiple functions: during long bone growth, FGF-18 is expressed in perichondrium and developing joints, and regulates bone formation by inhibiting chondrocyte proliferation and differentiation; FGF-18 knock-out mice survive embryonic development, but exhibit skeletal abnormalities and die in the early neonatal period. FGF-18 also induces ectopic cartilage

formation in the lung, and alters the morphology of the pulmonary

mesenchyma.

Protein Information

Name Fgf18

Function Plays an important role in the regulation of cell proliferation, cell

differentiation and cell migration. Required for normal ossification and bone development. Stimulates hepatic and intestinal proliferation (By similarity).

Cellular Location Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.