

Shh (C25II)

Catalog # PVGS1281

Product Information

Primary Accession Species	Q62226 Mouse
Sequence	Cys25-Gly198 (Cys25Ile-Ile)
Purity	> 95% as analyzed by SDS-PAGE > 95% as analyzed by HPLC
Endotoxin Level	
Expression System	E. coli
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O up to 100 µg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	20423
Other Names	Sonic hedgehog protein, SHH, 3.1.-.-, HHG-1, Shh unprocessed N-terminal signaling and C-terminal autoprocessing domains, ShhNC, Sonic hedgehog protein N-product, ShhN, Shh N-terminal processed signaling domains, ShhNp, Sonic hedgehog protein 19 kDa product, Shh {ECO:0000312 MGI:MGI:98297}, Hhg1
Target Background	Sonic Hedgehog (Shh) is a member of the Hedgehog (Hh) family of highly conserved proteins which are widely represented throughout the animal kingdom. In mammal, there are three related Hh proteins, Sonic (Shh), Desert (Dhh) and Indian (Ihh). They share a high degree of amino-acid sequence identity (e.g., Shh and Ihh are 93% identical). Sonic Hedgehog plays a role in cell growth, cell specialization, and the normal shaping (patterning) of the body. Shh is also important for development of the brain and spinal cord (central nervous system), eyes, limbs, and many other parts of the body.

Protein Information

Name	Shh {ECO:0000312 MGI:MGI:98297}
Synonyms	Hhg1
Function	[Sonic hedgehog protein]: The C-terminal part of the sonic hedgehog protein precursor displays an autoproteolysis and a cholesterol transferase activity (PubMed: 7736596 , PubMed: 7891723 , PubMed: 8824192). Both activities result in the cleavage of the full- length protein into two parts (ShhN and ShhC) followed by the covalent attachment of a cholesterol moiety to the C-terminal of the newly generated ShhN (PubMed: 8824192). Both activities occur in the reticulum endoplasmic (PubMed: 21357747). Once cleaved, ShhC is degraded in the endoplasmic reticulum (PubMed: 21357747).
Cellular Location	[Sonic hedgehog protein]: Endoplasmic reticulum membrane {ECO:0000250 UniProtKB:Q15465}. Golgi apparatus membrane {ECO:0000250 UniProtKB:Q15465}. Note=Co-localizes with HHAT in the ER and Golgi membrane. {ECO:0000250 UniProtKB:Q15465}
Tissue Location	Expressed in a number of embryonic tissues including the notochord, ventral neural tube, floor plate, lung bud, zone of polarizing activity and posterior distal mesenchyme of limbs In the adult, expressed in lung and neural retina

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.